

The Effect of Acupuncture on Telomere Length and Sperm Quality: A Research Synthesis

By Romy Simone Schwartz, L.Ac.

A Capstone Project

Submitted in partial fulfillment of the requirements for the degree Doctor of

Acupuncture and Oriental Medicine


Yo San University

Los Angeles, California


March 2017

**Approval Signatures Page**

This Capstone Project has been reviewed and approved for acceptance in fulfillment of  
DAOM Research Reporting by:

  
\_\_\_\_\_  
Rosaleen Ostrick, L.Ac.  
Capstone Project Advisor

*April 30, 2017*

  
\_\_\_\_\_  
Daoshing Ni, DOM, LAc, PhD  
Integrative Reproductive Medicine Specialty Chair

*April 30, 2017*

  
\_\_\_\_\_  
Lorraine Crampton, DAOM, LAc,  
Doctoral Program Dean      Date

*April 30, 2017*

### Abstract

Male infertility accounts for roughly 33% of the main causes of infertility among couples and is a major concern surrounding pregnancy outcomes and live birth rates. Current treatments are invasive and costly, and significant numbers of new findings and research suggest that these current forms of treatment such as Assisted reproductive technology (ART), Intra-cytoplasmic sperm injection (ICSI) and In vitro fertilization (IVF) have been shown to affect the fetus and cause problems later in life for those children. Acupuncture, which has been around for 5,000 years, holds promise as a treatment for sperm quality. Studies have shown that males have an increase in sperm count as well as faster movement of sperm with acupuncture treatment. Stress is a widely known factor that may affect male infertility because it relates to the aging process and overall decrease in general health and vitality. Telomeres, the end caps of chromosomes, are a general marker for a person's health, and their measurement may provide into the effects of stress on male infertility. Telomere length and telomerase levels are an important area of focus in medicine because studies are showing that measuring telomeres can show how a person is aging and also possible relationships between telomere length and cancer risk. The purpose of this study is to identify whether there is a correlation between the effects of acupuncture on telomere length, improvement in sperm quality and the ability of longer telomeres to improve sperm quality. This study also identifies the role of stress on factors related to male infertility and telomeres as well as how acupuncture may be shown to decrease stress. **Methods:** PubMed, Google Scholar, Yo San Library, EBSCO host databases and published books were explored for search terms including acupuncture, telomere length, telomerase levels, stress, male infertility and sperm quality. **Results:** Research revealed that male infertility and sperm quality are negatively affected by stress and acupuncture has been shown to minimize stress. In addition,

telomere length and telomerase were both shown to be inversely proportional to stress as well as sperm quality.

### **Acknowledgments**

I would first and foremost like to thank my sweetheart. He is always available to listen to me, nurture me, offer advice and just be there for me in everything that I may need. I feel blessed to have had him in my life for all of these years. Next, I would like to thank my father, whose unconditional love and support has always helped guide me through life. His compassion, understanding, courage and loving nature have all been blessings. He is a constant presence in my life through the joys, the challenges, and the ebbs and flow of life. I would also like to thank Rosaleen Ostrick, Lac, my capstone advisor. She really went above and beyond what was called for in helping me. I feel honored to have been lucky enough to get her as my advisor. Her expertise and kindness were gracious. I give thanks to Juliana Baldwin. There are some key professors, whose support and expertise in the field of my research has helped tremendously. Thank you to Dr. Jiling Hu, LAc, OMD, Dr. Hua Bing Wen, Lac, OMD, Denise Wiesner, LAc, Dr. Hong Yun Jin, LAc, OMD, Dr. Theodric Hendrix, MD, Dr. Catherine DeUgarte, MD, and Dr. Nurit Winkler, MD. I feel blessed and inspired to know each of you. Thanks to Dr. Andrea Murchison for her countless hours of help and support from the beginnings of formulating a Capstone subject matter. Also, thanks to Dr. Laraine Crampton who was able to pick up from where Andrea left off, while offering much needed support and advise. Last, but not least, thanks to my fellow cohort. We were a small but independent bunch of girls. You ladies helped make these two years a joy and pleasure to experience this doctoral program with.

## Table of Contents

<b>Approval Signature Page .....</b>	<b>2</b>
<b>Abstract.....</b>	<b>3</b>
<b>Acknowledgments .....</b>	<b>4</b>
<b>Chapter 1: Introduction .....</b>	<b>8</b>
<b>Male Infertility .....</b>	<b>8</b>
<b>Definition of Terms .....</b>	<b>8</b>
<b>The Scope of the Problem.....</b>	<b>8</b>
<b>Epidemiology .....</b>	<b>10</b>
<b>Known Causes of Male Unproductiveness .....</b>	<b>10</b>
<b>Diagnostic Testing for Male Infertility .....</b>	<b>12</b>
<b>Treatment Options for Male Infertility .....</b>	<b>13</b>
<b>Shortcomings of Current Male Infertility Treatments.....</b>	<b>14</b>
<b>Genetic Causes of Infertility .....</b>	<b>15</b>
<b>DNA Fragmentation .....</b>	<b>15</b>
<b>Telomere Length Affecting Male Infertility .....</b>	<b>17</b>
<b>What are Telomeres and Telomerase? .....</b>	<b>17</b>
<b>Telomere Length and Telomerase Levels in Sex (Germ) Cells V. in Regular (Somatic) Cells and the Significance of Each .....</b>	<b>19</b>
<b>Telomeres, Telomerase and Cancer .....</b>	<b>20</b>
<b>Male Infertility Correlation to Telomeres .....</b>	<b>21</b>

<b>Acupuncture and Male Infertility .....</b>	<b>21</b>
<b>Stress .....</b>	<b>23</b>
<b>How Stress Relates to Telomere Length and Telomerase Levels .....</b>	<b>23</b>
<b>How Stress Relates to Male Infertility .....</b>	<b>23</b>
<b>The Effect of Acupuncture on Stress .....</b>	<b>24</b>
<b>Statement of Research Question and Hypothesis .....</b>	<b>25</b>
<b>Definition of Key Terms and Glossary of Abbreviations .....</b>	<b>26</b>
<b>Chapter 2: Literature Review .....</b>	<b>31</b>
<b>Impact of Telomere Length on Sperm Quality .....</b>	<b>31</b>
<b>Effect of Acupuncture on Sperm Quality .....</b>	<b>34</b>
<b>Impact of Acupuncture on Telomere Length .....</b>	<b>36</b>
<b>Telomere Shortening Associated With Increased Risk of Cancer .....</b>	<b>37</b>
<b>Impact of Acupuncture on Stress .....</b>	<b>38</b>
<b>Chapter 3: Methodology .....</b>	<b>40</b>
<b>Designation of Methodology .....</b>	<b>41</b>
<b>Search Strategy .....</b>	<b>41</b>
<b>Inclusion and Exclusion Criteria .....</b>	<b>42</b>
<b>Data Collection .....</b>	<b>42</b>
<b>Human Subjects Ethical Consideration .....</b>	<b>43</b>
<b>Chapter 4: Results .....</b>	<b>43</b>
<b>Overview .....</b>	<b>43</b>
<b>Impact of Acupuncture on Telomere Length .....</b>	<b>44</b>
<b>Impact of Telomere Length on Sperm Quality .....</b>	<b>47</b>
<b>Impact of Acupuncture on Sperm Quality .....</b>	<b>51</b>
<b>Impact of Stress on Telomere Length .....</b>	<b>53</b>

**Impact of Stress on Male Fertility .....54**

**Chapter 5: Discussion .....57**

**Summary of Findings .....57**

**Study Limitations.....58**

**Theory of Implication .....58**

**Recommendations for Additional Studies .....58**

**Conclusion .....59**

**References:.....59**

**Appendix 1: Table 4.....76**

**Appendix 2: Table 5.....77**

**Appendix 3: IRB Proposal .....78**

**Appendix 4: Curriculum Vitae .....79**

## **Chapter 1: Introduction**

### **Male Infertility**

#### **Definition of terms**

Dohle et al., 2004, defined infertility as the inability of a sexually active, non-contracepting couple to achieve pregnancy within one year. The male partner can be evaluated for infertility using a laboratory evaluation of semen. According to Gnoth et al., 2005, a common definition of infertility is extremely useful for the suitable management of infertility. The term “infertility” may be used interchangeably with “sterility” with both resulting in only occasional occurrence of spontaneous pregnancies (Gnoth et al., 2005). The major factor affecting the individual spontaneous pregnancy prospect is the amount of time of undesirable non-conception, which determines the grading of infertility (Gnoth et al., 2005).

#### **The scope of the problem**

Dohle et al., 2004, state about 25% of couples do not achieve pregnancy within one year, of whom 15% of whom seek medical treatment for infertility and less than 5% of whom remain unwillingly childless. Schaeffer & Hotaling (2016) conceded that if just 15% of couples are infertile, with half due to male factor, then 7% of males have some degree of male factor infertility. Therefore, ten million men in this cohort aged 18–50

translates into 700,000 men with male infertility. When a man is infertile, difficulties are often related to sperm count or sperm quality. According to Gnoth et al., 2005, symptoms of male infertility include abnormally shaped sperm, sperm with poor motility, poor sperm quality, sperm that cannot attach the head to the egg and sperm that cannot penetrate the egg.

The quality of a man's sperm decreases with age (Gnoth et al., 2005). As a man ages, it takes longer for his partner to get pregnant (Gnoth et al., 2005). There's also an increased risk of not conceiving at all. The risk of miscarriage is higher if the father is over 45 (Neto, Najari & Goldstein, 2016). The children of older fathers are at greater risk of autism, mental health problems and learning difficulties (Neto, Najari & Goldstein, 2016). The average time to pregnancy if a man is under 25 is just over four and a half months but nearly two years if a man is over 40 (Neto, Najari & Goldstein, 2016). There is a five-fold increase in time to pregnancy if the male partner is aged over 45 years (Neto, Najari & Goldstein, 2016). For couples having In vitro fertilization (IVF), the risk of not having a baby is more than five times higher if the male partner is aged 41 or older (Taylor, 2003). The volume of a man's semen and sperm decrease continually between the ages of 20 and 80 (Taylor, 2003). The risk of miscarriage is twice as high for women whose male partner is aged over 45 than for those whose partners are under 25 (Taylor, 2003). Children with fathers aged 40 or older are more than five times as likely to have an autism spectrum disorder than children fathered by men aged under 30 (Taylor, 2003).

## **Epidemiology**

Ashok (2015) found that the male factor infertility ranged from 20% to 70% and that the percentage of infertile men ranged from 2.5% to 12%. Infertility rates were highest in Africa and Central/Eastern Europe. Rates of male infertility in North America, Australia, and Central and Eastern Europe varied from 4.5-6%, 9%, and 8-12%, respectively (Ashok, 2015). According to the results, at least 30 million men worldwide are infertile with the highest rates in Africa and Eastern Europe (Ashok, 2015).

## **Known causes of male infertility**

Gnoth et al, 2005, describe causes of male infertility in detail (See Table 1). Sperm production problems, blockage of sperm transport, erection and ejaculation problems, hormonal problems and sperm antibodies are the major known causes of male infertility (Gnoth et al., 2005). More than 90% of male infertility cases are due to low sperm counts, poor sperm quality, or both. (Jain, 2006). The remaining cases of male infertility can be caused by a range of conditions including anatomical problems, hormonal imbalances, and genetic defects (Jain, 2006). There are many causes of male infertility, which makes diagnosis difficult.

**Table 1: Known Causes of Male infertility (Gnoth et al., 2005)**

Sperm production problems	<ul style="list-style-type: none"> <li>• Chromosomal or genetic causes</li> <li>• Undescended testes at birth</li> <li>• Infections</li> <li>• Torsion (twisting of testis in scrotum)</li> <li>• Varicocele (varicose veins of the testes)</li> </ul>
---------------------------	--

	<ul style="list-style-type: none"> <li>• Medicines and chemicals</li> <li>• Radiation damage</li> <li>• Unknown cause</li> </ul>
Blockage of sperm transport	<ul style="list-style-type: none"> <li>• Infections</li> <li>• Prostate-related problems</li> <li>• Absence of vas deferens</li> <li>• Vasectomy</li> </ul>
Sexual problems (erection and ejaculation problems)	<ul style="list-style-type: none"> <li>• Retrograde and premature ejaculation</li> <li>• Failure of ejaculation</li> <li>• Erectile dysfunction</li> <li>• Infrequent intercourse</li> <li>• Spinal cord injury</li> <li>• Prostate surgery</li> <li>• Damage to nerves</li> <li>• Some medicines</li> </ul>
Hormonal problems	<ul style="list-style-type: none"> <li>• Pituitary tumors</li> <li>• Congenital lack of Luteinizing hormone (LH)/ Follicle-stimulating hormone (FSH) as a pituitary problem from birth</li> <li>• Anabolic (androgenic) steroid abuse</li> </ul>
Sperm antibodies	<ul style="list-style-type: none"> <li>• Vasectomy</li> <li>• Injury or infection in the epididymis</li> <li>• Unknown cause</li> </ul>

### **Diagnostic testing for male infertility**

Taylor (2003) states that diagnosing male infertility problems usually involves the following:

- General physical examination and medical history, including sexual habits and sexual development during puberty.
- Semen analysis; semen is sent to a laboratory to measure the number of sperm present and to look for any abnormalities in the shape (morphology) and movement (motility) of the sperm, or any infections.
- Since sperm counts often fluctuate significantly from one specimen to the next, semen analyses are generally done over a period of time for the most accurate results.

Table 2 details frequent diagnostic tests used for male infertility.

**Table 2: Diagnostic Tests Commonly Used for Male Infertility (Taylor (2003))**

<ul style="list-style-type: none"> <li>• Scrotal ultrasound to look for varicocele or other testicular problems.</li> </ul>
<ul style="list-style-type: none"> <li>• Hormone testing, since the pituitary gland, hypothalamus, testosterone and testicles are involved in sexual development and sperm production.</li> </ul>
<ul style="list-style-type: none"> <li>• Post-ejaculation urinalysis; sperm can travel backward into the bladder instead of out the penis during ejaculation (retrograde ejaculation).</li> </ul>
<ul style="list-style-type: none"> <li>• Genetic tests; usually for low sperm concentration. A blood test can test for changes in the Y chromosome, signs of a genetic abnormality and</li> </ul>

diagnose various congenital or inherited syndromes.
<ul style="list-style-type: none"> <li>• Testicular biopsy (by removing samples from the testicle with a needle) can determine if sperm production is normal, which then differentiates from a blockage or another possible problem with sperm transport</li> </ul>
<ul style="list-style-type: none"> <li>• Specialized sperm function tests to check how well the sperm survives after ejaculation, how well they can penetrate an egg, and whether there's any problem attaching to the egg.</li> </ul>

### **Treatment options for male infertility**

Available treatment options available for male fertility problems include surgery, medication, hormone treatments, use of donor sperm and various Assisted Reproductive technologies (ART) procedures (Taylor, 2003). The most common ART procedures currently used are In vitro fertilization (IVF), Intrauterine insemination (IUI) and Intra-cytoplasmic sperm injection (ICSI), which are defined in the definition of key terms section. Table 3 provides detailed treatment options for infertile males.

**Table 3: Treatments for Male Infertility (Taylor (2003))**

<ul style="list-style-type: none"> <li>• Assisted reproductive technology (ART) treatments, such as Intrauterine insemination (IUI), In vitro fertilization (IVF), or Intra-cytoplasmic sperm injection (ICSI: a single sperm is injected directly into an egg),</li> </ul>
<ul style="list-style-type: none"> <li>• A varicocele can often be surgically corrected, or an obstructed vas deferens repaired. Prior vasectomies can be reversed.</li> </ul>

<ul style="list-style-type: none"><li>• Treating infections of the reproductive tract with antibiotic therapy.</li></ul>
<ul style="list-style-type: none"><li>• Treatments for sexual intercourse problems, such as medication or counseling for erectile dysfunction or premature ejaculation.</li></ul>
<ul style="list-style-type: none"><li>• Hormone treatments and medications, such as injections of human chorionic gonadotropin and or recombinant human follicle stimulating hormone.</li></ul>

### **Shortcomings of current male infertility treatments**

Current treatments for male infertility have serious shortcomings (Sylvest et al. 2016). Intrusiveness or complications of surgeries, for example, surgery to remove a varicocele, may include pain, blood clots, heart disease risk, prostate cancer and dementia risks (Sylvest et al., 2016).

Dohle et al., 2004 states that In vitro fertilization (IVF) is a method used by reproductive endocrinologists to bypass many of the male factor sperm issues. However, IVF can sometimes create complications for the unborn child (Dohle et al., 2004). According to Dohle et al., 2004, more than 12 scientific papers have been published in the past year suggesting a connection between fertility treatments and low birth weight infants or disorders such as hypospadias (penis opening on back instead of front), heart malformations, chromosomal abnormalities or other major birth defects. In 2008, the Centers for Disease Control (CDC) noted that babies conceived with IVF had a slightly increased risk of several birth defects, including a hole in the heart, cleft palate, improperly developed esophagus and malformed rectum. There may be some abnormal

gene expression associated with IVF, which increases the chance of genetic disorders (Dohle et al., 2004). Advanced paternal age using IVF has been linked to autism, Marfan and Apert syndromes and other problems (Dohle et al., 2004).

### **Genetic causes of infertility**

Infertility is a broad term used to define a range of different phenotypes. According to Shah et al (2003), “The genetic causes of infertility are varied and include chromosomal abnormalities, single gene disorders and phenotypes with multifactorial inheritance”. The study also mentions that “specific genotypes and karyotypes have been associated with infertility and studies of specific genes in humans and model systems shed light on the nature of the polygenic and multifactorial basis of infertility” (p.17).

### **DNA fragmentation**

DNA fragmentation may predict the success of ART. ICSI may be used when there is high DNA fragmentation of sperm (Najari et al., 2016). Rodriguez et al (2005), states, “sperm chromatin abnormalities or DNA damage can impact human male fertility” (p.843). Evenson et al (2002) relates to this idea in stating that “sperm cells with sizable DNA breakage are generally present in human ejaculate, but infertile males typically have a higher proportion of sperm cells with DNA fragmentation” (p.36). DNA fragmentation is the separation or breaking of DNA strands into pieces (Carrell et al., 2009). The genetic integrity of the spermatozoon is essential for normal embryo development (Evenson et al., 2002). A high level of DNA fragmentation in sperm cells may represent a cause of male infertility that conventional examinations - sperm

concentration, motility analysis, and morphology assessment - cannot detect (Evenson et al., 2002). Regardless of the assisted reproductive technology used, elevated levels of DNA fragmentation above the critical threshold will significantly compromise the possibility of a successful pregnancy (Niederberger, 2016). A DNA fragmentation test is an effective method for measuring thousands of sperm in an ejaculate (Carrell et al., 2009). Sperm are stained with a fluorescent probe that interacts with the DNA molecule (Niederberger, 2016). The fluorescence signal changes when the DNA is fragmented, and this is monitored using a flow cytometer (Niederberger, 2016). Carrell et al., 2009, recommends that this test be done on all infertile males.

**Table 4: DNA Fragmentation Index (DFI): Percentage of Sperm Cells Containing Damaged DNA (Niederberger, 2016).**

<ul style="list-style-type: none"> <li>• <math>\leq 15\%</math> DFI = excellent to good sperm DNA integrity (IUI), In vitro fertilization</li> </ul>
<ul style="list-style-type: none"> <li>• <math>&gt; 15</math> to <math>&lt; 25\%</math> DFI = good to fair sperm DNA integrity</li> </ul>
<ul style="list-style-type: none"> <li>• <math>&gt; 25</math> to <math>&lt; 50\%</math> DFI = fair to poor sperm DNA integrity</li> </ul>
<ul style="list-style-type: none"> <li>• <math>\geq 50\%</math> DFI = very poor sperm DNA integrity</li> </ul>

## **Telomere Length Affecting Male Infertility**

### **What are telomeres and telomerase?**

Telomeres are the end caps of chromosomes (Rodriguez et al., 2005). They ordinarily shorten with each round of DNA replication, which ultimately leads to the inability of cells to divide, leading to cell death, according to Rodriguez et al., 2005. Telomeres are important because they protect the ends of our chromosomes, much like the plastic tip on shoelaces (Fajkus, Dvorácková & Sýkorová, 2008). If the telomeres were not there, our chromosomes may end up sticking to other chromosomes (Fajkus, Dvorácková & Sýkorová, 2008). Since chromosomes are shortened during each replication, the only part of the chromosome that is lost is the telomere, and the DNA is left undamaged (Fajkus, Dvorácková & Sýkorová, 2008). Without telomeres, important DNA would be lost every time a cell divides which would eventually lead to the loss of entire genes (Fajkus, Dvorácková & Sýkorová, 2008).

The chromosome telomere is a series of 5'-TTAGGG-3' repeats found at the terminal region of a chromosome (Rodriguez et al., 2005). According to Rodriguez et al (2005), “This repeat sequence plays a vital role in DNA replication and in the protection of the chromosome by inhibiting chromosomal fusions, and in chromosomal localization in the nucleus” (p.844). The length of the telomere varies between chromosomes and is shortened with every cell division by 50 to 200 base pairs (Rodriguez et al., 2005). Telomeres consist of the same sequence of bases repeated about 3,000 times and can reach up to 15,000 base pairs in length (Rodriguez, 2007). When the telomere becomes too short, the chromosome reaches a ‘critical length’ and can no longer be replicated,

which triggers the cell to die, a process called apoptosis, also known as cell death (Rodriguez, 2007). Rodriguez et al (2005) state, “All cells with a nucleus contain chromosomes and telomeres and exclude mature red blood cells and platelets” (p.844).

Telomeres ordinarily shrink by 1% annually, from birth to death. The telomeres of people with unhealthy habits have increased shrinkage (Rodriguez et al., 2005). Those of people with healthy habits and genes shrink at a slower rate, therefore helping these people to possibly live much longer (Rodriguez et al., 2005). Ultimately, telomere health is a significant determinant of health and longevity (Shealy (2016). Rejuvenation or regrowth of telomeres is, therefore, a crucial key to longevity and health according to Shealy (2016).

Table 5 provides the testing methods to measure telomere length. Each test analyzes either DNA, metaphase chromosomes or interphase nuclei (Montpetit et al., 2009).

**Table 5: Telomere Length Testing Methods (Montpetit et al., 2009)**

• HT Q-FISH: High throughput quantitative fluorescence in situ hybridization
• MMqPCR: Monochrome multiplex quantitative polymerase chain reaction
• qPCR: Quantitative polymerase chain reaction
• STELA: Single Telomere Length Analysis
• TRF: Terminal restriction fragment.
• PRINS: Primed in situ subtype of Q-FISH
• Q-FISH: Quantitative fluorescence in situ hybridization

Telomerase is an enzyme (reverse transcriptase) with protein + RNA subunit, that adds nucleotides to telomeres, prevents telomere shortening and repeated cell death (Fajkus, Dvorácková & Sýkorová, 2008). According to Fajkus, Dvorácková & Sýkorová (2008), “Telomerase is expressed by germ cells, early embryonic cells and stem cells, but not expressed by most somatic cells” (p.273).

The action of telomerase allows cells to keep multiplying and avoid aging (Fajkus, Dvorácková & Sýkorová, 2008). According to Fajkus, Dvorácková & Sýkorová (2008), “Because the lost sequences of the telomeres at each replication cycle can be resynthesized by telomerase, the enzyme has essential functions for cell immortalization” (p.275). Telomerase is important because it reverses telomere shortening and allows each generation to replace lost DNA, which allows cells to divide without ever reaching a limit (Fajkus, Dvorácková & Sýkorová, 2008).

### **Telomere length and telomerase levels in sex (germ) cells v. in regular (somatic) cells and the significance of each**

Chan & Blackburn (2004) state, “Telomere length has been demonstrated to both reflect and limit the replicative lifespan of normal somatic cells, any cell of a living organism other than the reproductive cells” (p.119). The replication of normal somatic cells is limited by telomere shortening, which additionally carries on with each round of cell division, resulting in the loss of 50–200 terminal base pairs of the telomere in humans both in vitro and in vivo (Lee et al., 2015). With ongoing rounds of DNA replication, the telomeres become shorter. However, telomerase can lengthen telomeres (Chan & Blackburn, 2004). Incomplete replication in the absence of telomerase results

in growing amounts of telomere shortening, which leads to the DNA damage and cell death (Chan & Blackburn, 2004). Therefore, proper telomere length and telomerase activity are necessary for replication of cells (Lee et al., 2015).

Telomerase is only found in very low concentrations in our somatic cells, whereas it is found in high levels in germ line cells (egg and sperm) and stem cells (Rudolph, 2007). Cells that divide rapidly, such as germ cells and stem cells, are among the few cell types in our bodies containing active telomerase (Rudolph, 2007).

### **Telomeres, telomerase and cancer**

Telomerase is also found in high levels in cancer cells (Shay et al, 2001). This enables cancer cells to be immortal and continue replicating themselves (Shay et al, 2001). According to Shay et al (2001), “If telomerase activity were switched off in cancer cells, their telomeres would shorten until they reached a critical length” (p.679). This would prevent the cancer cells from dividing uncontrollably to form tumors (Shay et al, 2001). However, blocking telomerase activity could affect cells where telomerase activity is important, such as sperm, eggs, platelets, and immune cells (Shay et al, 2001). Shay et al (2001) states, “Since telomerase activity in somatic cells is very low, these cells would therefore be largely unaffected by anti-telomerase therapy” (p.683).

Mu & Wei (2002) found that, “Telomerase activity may be detected in a variety of normal and malignant cell types by using a sensitive polymerase chain reaction-based method, known as the telomerase repeat amplification protocol (TRAP) assay” (p. 4).

Wesch, Burlock & Gooding, 2016 reported that telomerase activity was present in almost

all tumor samples but could not be detected in normal tissues, except for germ cells in the ovary and testis.

Telomerase activity is negative in human normal somatic cells but can be detected in most tumor cells (Mu & Wei 2002). Telomerase inhibitors might be useful in selectively killing cancer cells while saving normal cells (Wesch, Burlock & Gooding, 2016). According to Mu & Wei (2002), “The desired effect of telomerase inhibition would be to shorten telomeres to critical lengths, causing replicative senescence and preferably cell death due to irreparable chromosome damage. These objectives are feasible because the majority of tumors have shorter telomeres and higher proliferation rates than normal replicative cell populations, thus inhibition of telomerase activity may only be necessary over a few cellular divisions to attain lethal erosion of a telomere” (p.6). In contrast, Kim et al., 1994, found that the longer telomeres of stem cells and germ line cells should not lose their chromosomal protective function, allowing rather than inhibiting telomerase to prevent any telomere shortening.

### **Male infertility correlations to telomeres**

Abnormal shortening of telomeres has been associated with male factor infertility (Inagaki, 2006). Sperm maintain longer average telomere length because telomerase can rebuild the sperm cells if they begin to shorten, but sperm telomere length varies among individual men and individual spermatozoa (Inagaki, 2006). Variation in sperm telomere length could result from the effects of oxidative stress, which shorten telomeres (Inagaki, 2006). According to Kalmbach et al (2013), “Compromised telomere length in sperm

may contribute to segregation errors, apoptosis, programmed cell death, with reduced sperm count, and reduced fertility” (p.25).

### **Acupuncture and Male Infertility**

Acupuncture involves the insertion of fine and thin needles into the skin at important points on the body (Pei et al., 2005). Traditional Chinese medicine explains acupuncture as a technique for balancing the flow of energy through pathways (meridians) in the body (Pei et al., 2005). By inserting needles into specific points along these meridians, the energy flow will go back to homeostasis, according to the Pei et al., 2005. Acupuncture is an optional/additional treatment modality that can be used to enhance male fertility (Pei et al., 2005). According to Pei et al (2005), “Acupuncture infertility treatment produces few or no side effects while performing similar functions as the drugs do by stimulating the hypothalamus to effectively balance the endocrine system and its hormones and to get to the root cause of male infertility” (p.143).

Chen et al (2011) states that “With acupuncture, sperm production and quality can be increased, mechanical blockages can be dissolved or reduced and hormonal factors can often be corrected by restoring balance to the entire body” (p.313). Sufficient energy and substances and unblocked semen pathway are all necessary for ideal sperm quality (Pei et al., 2005). Three organ systems that can be addressed with acupuncture treatments are 1.) Kidney for growth and development, 2.) Liver for free flow of energy and 3.) Spleen for energy, sleep and sperm production (Chen et al., 2011). According to Sklar (2016), one of the ways acupuncture increases fertility is by reducing stress. Sherman et al (1997) states, “Acupuncture infertility treatment counters the effects of stress and cortisol by releasing endorphins in the brain, creating

a calming effect” (p.159). Acupuncture also increases fertility by strengthening the immune system, which can play an important role in conception (Sherman et al., 1997).

## **Stress**

Nargund (2015) states, “Psychological stress can be defined as any uncomfortable 'emotional experience' accompanied by predictable biochemical, physiological and behavioral changes or responses” (p.373). When people are under stress, the hormone cortisol is released in the brain (Mayorga-Torres et al., 2016). According to Nargund (2015), “This alters the brain's neurochemical balance, changing hormone levels and disrupting the pituitary balance” (p.381).

### **How stress relates to telomere length and telomerase levels**

Stress has been shown to decrease telomere length (Epel et al., 2004). Psychological stress is significantly associated with higher oxidative stress and shorter telomere length, which are known determinants of cell senescence and longevity (Zalli et al., 2014). According to Zalli et al (2014), “Higher levels of perceived stress decrease telomere length on average by the equivalent of at least one decade of additional aging compared to low stress” (p.4520).

### **How stress relates to male infertility**

Stress is a major underlying factor in infertility (Reynolds, 2016). Because of the delicate balance between the hypothalamus, pituitary, and reproductive glands, stress can alter sperm counts, motility, and cause impotence (Mayorga-Torres et al., 2016).

Emotional stress plays a detrimental role on fertility (Nargund, 2015). According to

Collodel (2008), “Data shows that, among sperm pathologies, necrosis and apoptosis were higher and the number of healthy sperm was significantly reduced in stressed men. The effects induced by stress also seem to include meiotic and structural alterations in sperm cells” (p.255).

McGrady (2009) looked at the effects of psychological stress on male fertility, showing that stress is associated with reduced paternity and abnormal semen parameters. Enough scientific evidence exists to suggest that psychological stress could severely affect spermatogenesis (McGrady, 2009). According to Collodel (2008), “The autonomic nervous system and the adrenal hormones participate in the classic stress response while also affecting the reproductive system. Evidence exists that mild-to-severe emotional stress depresses testosterone and perhaps interferes with spermatogenesis in the human male” (p.259).

### **The effect of acupuncture on stress**

Pilkington et al (2010), show that “studies on the endorphins released by acupuncture can raise the amount of white blood cells, T-cells and anti-bodies, which increase the body's level of immunity” (p.92). More recent work by Chao et al (1999) supports the concept that “Acupuncture activates the endogenous opioids in the central nervous system, which, in turn, inhibit central sympathetic neural outflow” (p.2128). According to Pomeranz (1988), “Evidence that acupuncture increases central nervous system opioids includes early work in a mouse model of pain, in which electro acupuncture at the large intestine 4(Li4) site produced analgesia” (p.228).

Yao & Thoren (1982) show “Experimental evidence in animal models of cardiovascular disease supports the concept that cardiovascular effects of acupuncture are mediated by inhibition of central sympathetic neural outflow, perhaps through release of endogenous opioids” (p.77). Eshkevari et al (2012) found that “Electronic acupuncture blocks the chronic, stress-induced elevations of the HPA axis hormones and the sympathetic NPY (Neuropeptide Y), a protein-coding gene pathway” (p.19). Sun et al (2005), report that “Experiments during acute stress show that electro-acupuncture (EA) at ST36 blocks chronic stimulation of the hypothalamic–pituitary–adrenal axis (HPA), and thus might reduce the physiological effects of the acute stress” (p.4963). Eshkevari et al (2015) also note that “Stress-induced circulating ACTH and corticosterone (CORT), as well as central CRH mRNA and protein were blunted when EA at acupoint stomach meridian point 36 (St36) was performed before exposure to chronic cold stress. This strongly suggests that EA at ST36 may be a viable therapeutic modality for chronic stress” (p.3649).

### **Statement of Research Question and Hypothesis**

The intent of this Capstone is to demonstrate the relationship between the effects of acupuncture on telomere length, improvement in sperm quality and the ability of longer telomeres to improve sperm quality. The role that increased psychological stress plays in this relationship is also discussed. The hypothesis of this researcher is that acupuncture can increase telomere length and sperm quality, and decreased stress levels in individuals can play an important role in positive outcomes.

## **Definition of Key Terms and Glossary of Abbreviations**

**ACTH: adrenocorticotrophic hormone** is a polypeptide tropic hormone produced and secreted by the anterior pituitary gland (Eshkevari et al, 2015).

**ART: assisted reproductive technology** is the technology used to achieve pregnancy in procedures such as fertility medication, artificial insemination, In vitro fertilization and surrogacy (Dohle et al., 2004).

**Asthenospermia:** is when less than 32% of spermatozoids move ((Siterman et al., 2000).

**Azoospermic:** is the complete lack of sperm in the ejaculate (Siterman et al., 2000).

**BDNF: brain-derived neurotrophic factor** provides instructions for making a protein found in the brain and spinal cord (Lin et al., 2015).

**CORT: corticosterone** is a 21-carbon steroid hormone of the corticosteroid type produced in the cortex of the adrenal glands (Eshkevari et al, 2015).

**CRH: corticotropin-releasing hormone** is a peptide hormone and neurotransmitter involved in the stress response (Eshkevari et al, 2015).

**DNA fragmentation:** is splitting the DNA into shorter pieces by endonucleolytic DNA cleavage at multiple sites (Rodriguez et al., 2005).

**EA: electro-acupuncture** is the application of a pulsating electrical current to acupuncture needles as a means of stimulating the acupoints (Gu et al, 2011).

**FRTA: free radical theory of aging** states that organism's age because cells accumulate free radical damage over time (Desai, Sabanegh, Kim & Agarwal, 2010).

**Genotype:** is the organism's full hereditary information (Esteves, 2013).

**Germ cell:** is any biological cell that gives rise to the gametes of an organism that reproduces sexually (Rodriguez et al., 2005).

**HPA: hypothalamic–pituitary–adrenal axis** is a complex set of direct influences and feedback interactions among three endocrine glands: the hypothalamus, the pituitary glands and the adrenal glands (Eshkevari et al, 2015).

**ICSI: intracytoplasmic sperm injection** is an assisted fertilization technique consisting of the microinjection of a single viable sperm into an extracted ovum (Siterman et al., 2000),

**Infertile phenotype:** is the physical characteristic(s) seen in an individual that renders him infertile along with genotype testing of chromosomes (Esteves, 2013).

**IUI: intrauterine insemination** is a fertility treatment that involves placing sperm inside a woman's uterus to facilitate fertilization (Siterman et al., 2000).

**IVF: in vitro fertilization** is the direct handling and manipulation of oocytes and sperm to achieve fertilization (Siterman et al., 2000).

**Karyotype:** is the number and appearance of chromosomes in the nucleus of a eukaryotic cell (Esteves, 2013).

**Knockout mice:** are laboratory mice in which researchers have inactivated, or "knocked out," an existing gene by replacing it or disrupting it with an artificial piece of DNA. Knocking out the activity of a gene provides valuable clues about what that gene normally does (Lin et al., 2015).

**Meiosis:** is the process of cell division by which reproductive cells (gametes) are formed (Desai, Sabanegh, Kim & Agarwal, 2010).

**NPY: neuropeptide y** is a 36-amino acid neuropeptide that acts as a neurotransmitter in the brain and in the autonomic nervous system of humans (Eshkevari et al, 2012).

**Oligozoospermia:** is when sperm concentration is less than 15 million/ml (Dieterle et al., 2009).

**OTA: oligo-teratoastheno-zoospermic** is a condition that includes low number of sperm, poor sperm movement and abnormal sperm shape (Siterman et al., 2000).

**Oxidative stress:** is a disturbance in the antioxidant balance, leading to potential damage (Aitken, Smith, Jobling, Baker & De Iuliis (2014).

**PESA: percutaneous epididymal sperm aspiration** is a procedure where a fine needle is passed into the tubes leading out of the testicles, or the testicles themselves, to retrieve sperm (Bidouee, Shamsa & Jalali (2011).

**Phenotype:** is an organism's observed property, such as morphology, development or behavior (Esteves, 2013).

**ROS: reactive oxygen species** are molecules or ions formed by the incomplete one-electron reduction of oxygen (Desai, Sabanegh, Kim & Agarwal (2010).

**Senescence:** is the condition or process of deterioration with age (Allsopp & Harley, 1995).

**Sham acupuncture therapy:** is essentially fake acupuncture, and is used as a control in scientific studies to determine whether or not the effects of a treatment are actually due to acupuncture treatment (Lin et al., 2015).

**Somatic cells:** are all cells in the body except germ cells, which are egg and sperm (Dohle et al., 2004).

**Spermatocyte:** is a male germ cell derived from spermatogonia (Dohle, et al. 2004).

**Spermatagonia:** are cells produced at an early stage in the formation of spermatozoa, formed in the wall of a seminiferous tubule and giving rise by mitosis to spermatocytes (Gnoth et al., 2005).

**Spermogram:** is a test where the male partner produces a sperm sample by masturbation into a sterile container (Bidouee, Shamsa & Jalali, 2011).

**Sperm morphology:** is the size and shape of sperm and is one factor that's examined as part of a semen analysis to evaluate male infertility (Gnoth et al., 2005).

**Sperm motility:** describes the ability of sperm to move properly through the female reproductive tract to reach the egg (Gnoth et al., 2005).

**Telomerase:** is an essential Ribonucleoprotein Reverse Transcriptase essential that adds telomeric DNA to the ends of eukaryotic chromosomes (Esteves, 2013).

**Telomere:** is a terminal section of a chromosome which has a specialized structure and which is involved in chromosomal replication and stability (Esteves, 2013).

**Teratozoospermia:** is abnormal sperm shape (Siterman et al., 2000).

**TH: tyrosine hydroxylase** is the rate-limiting enzyme in catecholamine synthesis (Benavides & Piccione, 2003).

**Upregulation:** is a positive regulatory effect on physiological processes at the molecular, cellular, or systemic level (Lin et al., 2015).

## Chapter 2: Review of Literature

### Impact of Telomere Length on Sperm Quality

Abnormal shortening of telomeres has been associated with male factor infertility (Esteves, 2013). The functions of telomeres (as stated in Chapter 1) include protection of the genetic information encoded on the chromosome, localizing the chromosomes in the nucleus and supporting DNA replication (Esteves, 2013).

Per Esteves (2013) study, “Telomeres have been targeted as potential candidates to explain some infertile phenotype characteristics in males” (p.177). Esteves’ study, a clinical appraisal of the genetic basis in unexplained male infertility, examined the relationship between genetics and unexplained male infertility. Esteves found that currently, few diagnostic tools are currently available for routine use and their usefulness is not yet completely clear.

Chromosomal defects are the most common genetic abnormalities in infertile males, accounting for 2.1-15.5% of cases. According to Esteves (2013), “Klinefelter syndrome, chromosome translocations, inversions and deletions fall in this category and the vast majority of affected individuals display severely compromised semen quality. Translocation carriers, however, may have varying sperm production phenotypes ranging from normal spermatogenesis

to inability to produce spermatogonia. Chromosomal abnormalities in somatic cells can be detected by karyotyping” (p.180).

Like chromosomal defects, gene mutations are usually related to severely abnormal sperm production phenotypes (Esteves, 2013). According to Esteves (2013), “Micro deletions in the Y chromosome, mutations in the cystic fibrosis gene and mutations of the androgen receptor gene are classic examples of genetic abnormalities in this disease category.

There is also a risk of unbalanced gamete production that results in an increased risk of spontaneous abortion and unbalanced offspring. The most relevant clinical aspect involves the carriers of translocations in chromosome 21 as they are at risk of producing a child with Down syndrome” (p.182).

Esteves (2013) states that “Sperm aneuploidy assessment in couples with unexplained infertility experiencing either repeated IVF failures or recurrent pregnancy loss can be performed by FISH while mutations and polymorphisms are identified by specific gene sequencing and mutational analysis methods” (p.182). Many advances are currently being made and the use of new genetic technology may hold the key to more accurately diagnosing and treating men with unexplained infertility (Esteves, 2013).

Desai, Sabanegh, Kim & Agarwal (2010) studied a free radical theory of aging and its implications in male infertility with regards to telomere shortening in humans. They studied the effects of mitochondrial generation of reactive oxygen species (ROS) and aging on human spermatozoa and seminal antioxidants and found that germ cell telomeres may be partially responsible for telomere shortening, decrease in sperm count, decline in testosterone concentration, and decline in motility with aging. They propose that ROS generation has a central role in the pathophysiology of age related decrease in male fertility.

Aitken, Smith, Jobling, Baker & De Iuliis (2014) studied oxidative stress and its impact on male reproductive health. Most of the discussion focused on the impact of oxidative stress at the level of gamete (Aitken, Smith, Jobling, Baker & De Iuliis, 2014). According to Aitken, Smith, Jobling, Baker & De Iuliis (2014), "If the oxidative stress is earlier in spermatogenesis, there are consequences for fertility and the health and well being of the offspring. Severe oxidative DNA damage in germ cells entering meiosis will simply precipitate an increase in cell death" (p.36).

An increase in telomere length has been shown to enhance sperm quality (Aitken, Smith, Jobling, Baker & De Iuliis, 2014). The impact of paternal aging on telomere length via aging is associated with oxidative stress in the germ line and increases telomerase activity and telomere length increase in spermatozoa (Aitken, Smith, Jobling, Baker & De Iuliis, 2014). According to Aitken, Smith, Jobling, Baker & De Iuliis (2014), "If the paternal germ line has experienced an oxidative stress post-meiotically, when telomerase can no longer increase (as is typically the case in infertile patients) then telomere length in the spermatozoa will be abnormally short and the implications for the health of ART offspring, potentially serious" (p.38). These studies showed how increased telomere length enhanced sperm quality.

Rodriguez and Herrera both used mice to study how telomere length affects male infertility. Rodriguez et al. (2005) studied the influence of excessive telomere shortening, a well-known trigger of sperm DNA fragmentation in late-generation knockout mice. Telomerase RNA Component (Terc) knockout mice are laboratory mice in which researchers have inactivated, or "knocked out," an existing gene by replacing it or disrupting it with an artificial piece of DNA. Knocking out the activity of a gene provides valuable clues about what that gene normally does (Lin et al., 2015). DNA fragmentation levels in sperm cells were measured from

wild type, early, and late generation telomerase null mice (Rodriguez et al, 2005). A minimum of 1,000 spermatozoa was assessed from each mouse (Rodriguez et al, 2005). Terc knockout mice had a six fold mean increase in the percentage of sperm cells with fragmented DNA. Lin et al (2015) study showed “critically shortened and highly eroded telomeres in Terc mice and found testicular atrophy and depletion of germ cells, possibly leading to chromosomal abnormalities. DNA fragmentation causes common infertility in males. This could genetically inactivate sperm cells with defective genomic constitution, preventing the development of offspring with an abnormal genetic background and thus resulting in decreased fertility” (p.36).

### **Effect of Acupuncture on Sperm Quality**

Research has shown that sperm quality can be positively influenced by acupuncture. Siterman et al. (2000), in a pilot study, showed a positive effect of acupuncture on sperm count, especially on men with such low sperm counts (or no sperm) that they would usually require a testicular biopsy to extract sperm for use in an IVF cycle. Semen samples of 20 males were taken. A light microscope and a scanning electron microscope then examined the spermatozoa before and a month after acupuncture treatments. Seven of the fifteen men with no sperm at all produced sperm detectable by the light microscope after a course of ten acupuncture treatments ( $p < 0.01$ ) i.e. enough sperm could be produced for ICSI to be performed without having a testicular biopsy. Specifically, a definite increase in sperm count was detected in the ejaculates of ten of the fifteen azoospermic patients and a marked but not significant improvement in the sperm counts of severely OTA (oligo-teratoastheno-zoospermic), low number of sperm, poor sperm movement and abnormal sperm shape, of males following acupuncture treatment.

Bidouee, Shamsa & Jalali (2011) presented a case study with single azoospermic male in whom spermatogenesis occurred who had sperm in his ejaculate after two courses of acupuncture. PESA (percutaneous epididymal sperm aspiration) using a fine needle to retrieve sperm and ICSI procedures had been done with failed results. Twenty acupuncture treatments twice a week using sterile disposable stainless steel needles ( $0.25 \times 25$  mm and  $0.25 \times 50$  mm) were administered for twenty-five minutes of needle retention to points for a deficiency of the kidneys (hormonal imbalance) and damp-heat syndromes (inflammation of the genital tract). Only twelve of these points were used at a time during each session. His next spermogram showed a sperm count of 10 million/mL, with 10% good and progressive motility and 60% normal shape, in the same lab. He then received a second course of acupuncture. His third spermogram, in the same lab, showed a sperm count of 18 million/mL, with 30% good and progressive motility and 60% normal sperm.

Siterman et al. (1997) studied a group of infertile men with abnormal semen analysis who were randomly divided into two groups; one group was given ten acupuncture treatments over five weeks, and the other group, no treatment. Significant improvements ( $p < 0.05$ ) were demonstrated in the acupuncture group compared to the control group; in particular, improved motility and morphology were shown. Sixteen acupuncture treated subfertility patients and 16 control untreated subfertility males were examined (Siterman et al, 2007).

Pei et al. (2005) studied a group of infertile men who had pathological semen analyses. They were treated with acupuncture twice a week for five weeks. A statistically significant increase after acupuncture in the percentage and number of sperm with no structural defects was demonstrated compared to the control group of patients who received no treatment. They concluded that male infertility patients could benefit from having acupuncture. Forty men with

idiopathic oligospermia, asthenospermia, or teratozoospermia were studied. Twenty-eight of the patients received acupuncture twice a week over a period of 5 weeks. The samples from the treatment group were randomized with semen samples from the 12 men in the untreated control group (Pei et al, 2005).

Dieterle et al. (2009) randomized fifty-seven patients who had extremely low sperm counts into acupuncture and placebo acupuncture groups. After receiving acupuncture twice weekly for six weeks motility of sperm (but not overall count) was found to increase significantly in the acupuncture group. The authors conclude that the results of the present study support the significance of acupuncture in male patients with severe oligoasthenozoospermia. More evidence with larger trials needs to be accumulated before the efficacy and effectiveness of acupuncture in male infertility can be evaluated.

Chen et al. (2011) studied the effect of acupuncture-moxibustion therapy on sperm quality in 118 infertility patients with sperm abnormality. The team found that levels of an important enzyme in the semen, seminal plasma acid phosphatase increased, as did the sperm motility after acupuncture, and were statistically significant ( $P<0.01$ ).

### **Impact of Acupuncture on Telomere Length**

Yoshiaki Omura performed a series of studies in 1998, 2007 and 2010 on the acupuncture point St 36 on humans to test its possible effects on telomere length. Omura, Shimotsura, Ooki, & Noguchi (1998), found that acupuncture on St.36 on apparently normal subjects increased telomere length up to a maximum of more than two times prior to the treatment, depending on the method of treatment, but increases were frequently between 60% to 100%. Subjects with unusually low telomere length often had chronic degenerative diseases, while those having

exceptionally high telomere length often had excellent physical conditions or mental acuity. (Omura, Shimotsura, Ooki, & Noguchi, 1998).

Lin et al. (2015) studied acupuncture's effect on telomerase response in the brain of mice using St 36. To investigate the telomerase response in the brain to acupuncture, brain-derived-neurotrophic factor (BDNF) levels were studied in three randomly assigned subgroups of mice, one receiving acupuncture (acupuncture subgroup), one-receiving sham acupuncture therapy (sham subgroup), and one-receiving no treatment (control subgroup). The mice were either telomerase-deficient (*Terc*) or normal, wild type (WT). The mice receiving acupuncture did so for four days at 30 minutes each at St 36. For the two sham groups, the sham point was set at a location approximately 3mm to the lateral side of the tail on the gluteus muscle following the same schedule. Acupuncture at St 36 for four days upregulated BDNF of the telomerase-deficient (*Terc*) mice, but that result was not seen in the normal, wild-type (WT) mice with normally functioning telomerase. The research team believed that using acupuncture increased telomerase activity through the activation of BDNF and its downstream, activating pathways in populations of patients who exhibit low telomerase activity, thus increasing telomere length (Lin et al., 2015).

### **Telomere shortening associated with increased risk of cancer**

Omura et al. (2007) studied an acupuncture point named "True St 36", a name coined by Omura, since the location varied slightly from the actual St 36 point location. Omura studied the cells of cancer patients and the results showed reduction in telomerase levels of these cancer cells, while increasing telomere length in the normal somatic cells, and in cells that did not exhibit cancer (Omura et al, 2007). Most human cancers have short telomeres and

express high levels of telomerase, whereas in most normal somatic tissues telomerase is absent (Shay et al, 2001). According to Shay et al (2001), “Tumors can arise from genetically instable cells with defective telomeres. Since telomere shortening is heavily associated with an increased risk of cancer during aging and chronic disease, the deficiency of telomere capping function adds to the induction of chromosomal instability and cancer initiation process” (p.683).

In their third study, Omura et al. (2010) studied the effect of acupuncture with electrical stimulation at St 36 on telomere length in breast cancer and Alzheimer’s. The results over the course of 10,20 and 30 minutes treatments showed telomere length had drastically increased in size in the normal somatic cells that were unaffected by the cancer, whereas cancer-related markers and abnormally high  $\beta$ -Amyloid levels in Alzheimer’s patients steadily decreased (Omura et al, 2010).

### **Impact of Acupuncture on Stress**

Sparrow & Golianu (2014) noted that there is evidence that acupuncture decreases the stress response in both human and animal subjects. This study measured the blood pressure of 10 hypertensive patients aged 33-72 and was a retrospective, uncontrolled case study. The researchers found that acupuncture can increase heart rate variability (HRV) in the short term (minutes to hours), which created a relaxed and low-stress environment. Sparrow & Golianu, (2014) state that “HRV has been linked to other accepted and precise physiologic markers of stress and inflammation pertinent to acupuncture treatment, one of them being telomere length, suggesting that the relaxed and calming HRV response may be applicable to clinical outcomes” (p.291).

Kondo & Kawamoto, 2014) found that acupuncture and moxibustion can improve the psychological relationship between doctors and patients, especially when it is disturbed by a “game”, a dysfunctional interpersonal interaction that is repeated unintentionally. Kondo & Kawamoto (2014) state that “Acupuncture and moxibustion are helpful in treating somatoform disorders based on fundamental meridian points that are considered to have effects on central, autonomic nervous, immune, metabolic, and endocrine systems to regulate the whole body balance” (p.7).

Nakamura et al. 2010, reported in a factor analysis of 197 patients that the attentive attitude of the therapist and proper touching by the therapist were significantly associated with a decrease in the subjective symptoms by acupuncture and moxibustion therapy, while the physical factors of the therapeutic stimulation were not. These results indicate that acupuncture and moxibustion therapy is a kind of psychotherapy, which may be of assistance in establishing the psychological therapist-patient relationship, rather than a physical therapy (Nakamura et al, 2010). It was recently reported that some of these meridian points have effects on the autonomic nervous system. (Kondo & Masazumi, 2014). For example, Ren 12 and ST 36 increase parasympathetic nerve activity (Liu, Xu & Chen 2012). In addition, some of the meridian points have opposite effects, which are contrary to other points. LI 11 and GB 21 increase sympathetic nerve activity (Knardahl, Elam, Olausson & Wallin, 1998 and Yook et al, 2009). Ngai & Jones 2013 found that UB 13 decreases sympathetic nerve activity.

Kondo & Masazumi (2014) note, “Since the acupuncture point combinations include meridian points with bi-directional regulation of the autonomic balance, they can be applied to various pathological conditions in a balanced manner” (p.7). Inoue and Yamauchi, 2008, Shea et al, 2008 and Tonhajzerova et al, 2010 showed that acupuncture reduced sympathetic and

parasympathetic nerve activities in people with both chronic anxiety disorders and severe depression. Minagawa et al, 2010, found that UB 32 reduces heart rate without having an effect on heart rate, which might be effective for patients with palpitations without heart rate variability abnormalities. Homma (1984) found that UB 10, UB 18, Ren 6, Ren 12, GB 20, DU 12, KD 6, LI 11 and St 36 have been shown to decrease anxiety. Park et al (2010) found that “Acupuncture for St 36 decreased anxiety-related behavior, the serum corticosterone level, and tyrosine hydroxylase-immunoreactive expression, the rate-limiting enzyme in catecholamine synthesis, of rats under immobilization stress” (p.113). Chen (1992) found that “UB18, Ren12, St 36, GB20, GB21 LR14, and St 36 have an anti-depressive effect. Electro-acupuncture at St 36 and GB20 reduced the Beck Depression Inventory scales in subjects with psychosomatic or psychiatric disorders such as fibromyalgia, irritable bowel syndrome, chronic fatigue syndrome, primary insomnia, and obsessive-compulsive disorder, probably by enhancing release of serotonin” (p.280). Eshkevari et al (2012) found that ST25 prevented chronic stress-induced increases in the sympathetic peptide, neuropeptide Y(NPY).

### **Chapter 3: Methodology**

The goal of this study was to explore whether telomere length affects sperm quality, whether acupuncture can increase sperm quality and whether acupuncture can therefore affect telomere length. This Capstone also sought to determine whether acupuncture’s affect on sperm quality might be due to acupuncture’s ability to reduce stress, which has been shown to affect sperm quality. This author’s hypothesis is that acupuncture increases telomere length, which results in higher sperm quality.

### **Designation of Methodology**

This study uses a systematic quantitative literature review focused on the following:

- Does telomere length affect sperm quality?
- Can acupuncture increase telomere length?
- Can acupuncture therefore affect sperm quality?
- Is acupuncture's affect on sperm quality due to its ability to reduce stress, or is there evidence that acupuncture's affect on sperm quality is independent of its ability to reduce stress?

A quantitative literature review was chosen as the methodology for this study. Pickering and Byrne (2013) define the benefits of systematic quantitative literature reviews for PhD candidates and other early career researchers. The review is systematic in terms of how papers are initially assessed for inclusion. By mapping the boundaries of the existing literature it is possible to identify gaps in the literature, determine where generalizations occur, and identify the limits of those generalizations (Pickering and Byrne, 2013). A hypothesis was developed, and then a systematic literature review was performed to compile and analyze information.

### **Search Strategy**

1. Online search of medical journals and peer-reviewed articles were performed through PubMed, NIH, Ebsco AltHealthWatch and Google Scholar.
2. Search terms used included: acupuncture and sperm, acupuncture and infertility, acupuncture and telomeres, acupuncture and telomerase, acupuncture and telomere

length, acupuncture and male infertility, acupuncture and sperm quality, telomeres and aging, acupuncture and sperm quality, acupuncture and sperm count, telomeres and sperm count, telomeres and male fertility, telomerase and male infertility, telomere length and sperm count, and telomeres and sperm quality.

3. Research for this study was collected from the United States, Japan, China, Brazil, Germany, Spain and Australia.
4. The articles used were all in English.
5. Of the 147 articles researched, 41 were used that fit the inclusion criteria.
6. Aside from articles, published books were used for research purposes.

### **Inclusion and Exclusion Criteria**

The study included research from 2000 or later. Human and animal studies were included. Articles in English were included.

Excluded from the study were non-peer-reviewed articles. Articles not in English were excluded. Articles older than 2000, were excluded, in order to focus on the most current research.

### **Data Collection**

A data abstraction form was used to extract relevant information for the current study. Criterion assessed included, name of article, date of publication, research method used and any outcomes attained in regards to each of the ensuing topics:

- A. How telomere length can affect sperm quality

- B. How acupuncture can affect telomere length
- C. How acupuncture can improve sperm quality
- D. How psychological stress affects telomere length and sperm quality

Once this data was collected, it was arranged into topics and sub-topics. These topics included acupuncture, telomeres and sperm quality. The sub-topics included: telomeres and telomerase, telomere length and telomere levels in germ and somatic cells, telomeres, telomerase and cancer, stress relating to telomere length and telomerase, stress relating to male infertility and the effects of acupuncture on stress. When studies with animals were used, the type of animal model used was specified.

### **Human Subjects Ethical Consideration**

This current study is a literature review synthesis. Therefore, there are no human subjects being used. In this case, no ethical considerations need to be taken into account. This research proposal was reviewed and approved by the Yo San University Institutional Review Board (IRB). A copy of the letter of approval is included.

## **Chapter 4: Results**

### **Overview**

This chapter will summarize the findings of the studies conducted on acupuncture, telomere length, sperm quality and stress. The studies were grouped into categories based on similar research article findings. These categories were: telomere length and sperm quality,

acupuncture and telomere length, acupuncture and sperm quality and stress affecting sperm quality and telomere length. A comparison of various acupuncture points was used, and animal or human studies and control groups were noted.

Thirty-eight original articles were reviewed that studied telomere length and acupuncture, acupuncture and male fertility and telomere length and male fertility. Upon further investigation, a question regarding what other factors may affect the outcomes of these studies became apparent. As a result, this author added the effect of stress on telomere length and the effect of stress on male fertility to her literature review. Ten additional original articles on stress and telomere length and stress and male fertility were reviewed as well. From the research, five groups were categorized into tables. These tables were compared and an analysis was presented based on the results of the findings.

### **Impact of Acupuncture on Telomere Length**

From the ten articles reviewed that addressed acupuncture's effects on telomere length, three directly showed that acupuncture positively affects telomere length. One of three of those articles also examined use of electrical stimulation on the acupuncture point. Omura, Shimotsura, Ooki & Noguchi (1998) found that acupuncture at St 36 on the cancer patients treated showed a decrease in telomere length in adenocarcinoma from 950 nanograms (ng) to 750 ng and a decrease in telomere length in small cell carcinoma from 770 ng to 600 ng. Normal subjects treated showed an increase in telomere length at least double and sometimes between 60-100%. The study conducted in 2007 by Omura, Chen, Lu, Shimotsura, Ohki & Duvvi, used "Press Needle Stimulation of Omura's St 36 repeated 200 times, 4 times a day to cancer patients, which reduced cancer cell telomere length by 600-1500ng and high Oncogene C-fos Ab2 and

Integrin  $\hat{I} \pm 5\hat{I}21$  of 100-700ng BDORT units to close to 1yg (=  $10\hat{a}^{24}$  g) BDORT units with a significant reduction of asbestos and mercury (Hg) from cancer cells, while markedly reduced normal cell telomere of 1yg was increased to optimally high amounts of 500 $\hat{A}$ -530ng BDORT units” (p.60). The significance of these numbers indicated that the needle stimulation was measured to have reduced cancer cell telomere length, while have increased the telomere length of normal cells.

Other researchers, such as Lin, et al (2015) also measured St 36 acupuncture point in mice to look at its effects on telomere length. Lin et al (2005) results found “e Terc-/- group showed down regulated hippocampal BDNF expression compared with the WT mice and acupuncture at St 36 for 4 days upregulated BDNF, TrkB, p75NTR, Akt, and ERK1/2 in the DG and hippocampus of the telomerase-deficient mice, but that result was not seen in the WT mice with normally functioning telomerase” (p.36). This research concluded that the use of acupuncture in pathologies associated with telomerase deficiency might provide some benefit by activation of brain-derived neurotrophic factor. Of the seven remaining articles, the results ended up being too heterogeneous to draw a conclusion.

**Table 1: Overview of Studies (Acupuncture and Telomere Length)**

1 <sup>st</sup> Author (year)	Subjects	Measured	Intervention	Acupuncture points used	Results
Omura (1998)	*30 normal subjects *1 cancer	Telomere length	Acupuncture	Omura's ST 36	Acupuncture increased telomere length in normal

	subject				cells and decreased length in cancer cells
Omura (2007)	Cancer patients: 28 y/o female 72 y/o female	Telomere length	Acupuncture	Omura's ST36	Acupuncture can decrease telomere length in cancer patients
Omura (2010)	*69 y/o female with cancer *79 y/o male with Alzheimer's	Telomere length	*TEM (Transcutaneous Electrical Stimulation ) *Acupuncture	Omura's ST 36	Increasing normal cell telomeres by Omura's ST36 stimulation by press-needle stimulation, and increasing normal cell telomeres by transcutaneous electrical stimulation of Omura's ST36
Lin (2015)	*Terc-telomerase deficient mice *WT-wild-type	Telomere length	Acupuncture	ST 36	The Terc-/- mice in all subgroups had downregulated hippocampal BDNF expression compared with that

					of the WT mice in all subgroups.
--	--	--	--	--	----------------------------------

### **Impact of Telomere Length on Sperm Quality**

From the twelve articles researched on telomere length and sperm quality, five showed a direct correlation between telomere shortening with decreased male fertility. Of those five articles, two were mice studies and three were human studies. Studies in mice have proposed that there is a safety mechanism which degenerates spermatocytes with diminished telomere length to prevent their maturation. Rodriguez et al (2005) found that “Terc knockout mice had a six-fold mean increase in the percentage of sperm cells with fragmented DNA, since these mice of G3 (generation 3) had critically short telomeres” (p.844). Herrera et al (1999) study showed that in different generations of mice, the average length of telomeres was significantly shorter in the 6<sup>th</sup> generation mice than the earlier generations. According to Herrera et al (1999), “Mice lacking telomerase activity can only be bred for approximately six generations due to decreased male fertility and to an increased embryonic lethality associated with a neural tube closure defect” (p.2956). Of the seven remaining articles, the results ended up being too heterogeneous to draw a conclusion.

Two of the human studies both examined free radical theory and its effects on sperm

quality. Desai et al (2009) examined “reactive oxygen species (ROS), important signaling molecules in an organism's regulation of metabolism and inflammation and male infertility.

Accumulation of ROS may be responsible for telomere shortening and subsequent decrease in sperm count, decline in testosterone concentration, and decline in motility with aging” (p.17).

Aitken et al (2013) found oxidative stress affecting the male with telomere length in the spermatozoa being abnormally short and the implications for the health of ART offspring, potentially serious. Emery & Carrell’s (2006) review showed that, “the telomere checkpoint is leaky in spermatocytes. Additionally, fertilization with telomerase null sperm causes severe defects in the resultant embryos” (p.139). These observations might indicate that these sperm are dysfunctional as a result of short telomeres in mature sperm (Emery & Carrell, 2006).

According to Emery & Carrell (2006), “The telomere plays a major role in chromosomal localization within the sperm nucleus and it is possible that abnormal localization could cause meiotic errors resulting in the presence of an abnormal number of chromosomes in the sperm” (p.140).

**Table 2: Overview of Studies (Telomere Length and Sperm Quality)**

<b>1<sup>st</sup> Author(year)</b>	<b>Subjects</b>	<b>Male factor</b>	<b>Measured</b>	<b>Results</b>
<b>Rodriguez(2005)</b>	<b>14 mice (WT ,G1 Terc, G3 Terc)</b>	<b>DNA sperm fragmentation</b>	<b>Telomere shortening</b>	<b>Critically short telomeres are associated with sperm DNA</b>

				<b>fragmentation</b>
<b>Herrera (1999)</b>	<b>18 G3 mice</b>	<b>Decreased fertility</b>	<b>Short telomeres</b>	<b>Increased male infertility associated with telomerase deficiency appear earlier in mice with short telomeres</b>
<b>Desai(2009)</b>	<b>Human male subjects</b>	<b>Decreased fertility</b>	<b>Telomere shortening</b>	<b>Increases ROS responsible for telomere shortening &amp; subsequent decrease in sperm count &amp; motility decline</b>

<b>Aitken(2013)</b>	<b>Human male subjects</b>	<b>Decreased fertility</b>	<b>Telomere shortening</b>	<b>Oxidative stress is a major pathological mechanism responsible for both male infertility and DNA damage in the germ line.</b>
<b>Emery(2006)</b>	<b>Human male subjects</b>	<b>Sperm abnormalities</b>	<b>Telomere length</b>	<b>Fertilization with telomerase null sperm causes severe defects in the resultant embryos.</b>

### **Impact of Acupuncture on Sperm Quality**

Of the sixteen articles reviewed on the impact of acupuncture on sperm quality, six directly correlated acupuncture treatments with increased sperm quality. Siterman et al (1997)

found that “patients exhibiting a low fertility potential due to reduced sperm activity may benefit from acupuncture treatment, with a fertility index increase significantly ( $p < \text{or} = .05$ ) following improvement in total functional sperm fraction, percentage of viability, and total motile spermatozoa” (p.156). Siterman et al (2000) performed another study, but this time with men with low to no sperm counts and found and found detectable sperm after the acupuncture ( $p < 0.01$ ). As a result, enough sperm could be produced for ICSI to be performed. Again, in 2009, Siterman et al, measured acupuncture results which improved sperm quality by decreasing the temperature of the testes. Siterman et al (2009) results found “Seventeen of the 34 patients with hyperthermia, all of whom had genital tract inflammation, had normal scrotal skin temperature; in 15 of these 17 patients, sperm count was increased” (p.205).

Pei et al (2005) found a general improvement of sperm quality, specifically in the ultra structural integrity of spermatozoa, after acupuncture treatment. In their case study, Bidouee et al (2011), treated a 31 year old male with a history of failed PESA + ICSI before acupuncture. The results showed he had a sperm count of 18 million/mL in his ejaculation, after two courses of acupuncture. His next spermogram showed a sperm count of 10 million/mL, with 10% good and progressive motility and 60% normal shape, in the same lab. Following this, he received a second course of acupuncture. His third spermogram, in the same lab, showed a sperm count of 18 million/mL, with 30% good and progressive motility and 60% normal sperm. In measuring severely low sperm count as well, Dieterle et al (2009) studied 57 patients who had extremely low sperm counts, to acupuncture and placebo acupuncture groups. After receiving acupuncture twice weekly for 6 weeks motility of sperm (but not overall count) was found to increase significantly.

**Table 3: Overview of Studies (Acupuncture and Sperm Quality)**

<b>1<sup>st</sup> author(date)</b>	<b>Subjects</b>	<b>Study/Control</b>	<b>Intervention</b>	<b>Measured</b>	<b>P-value/ Results</b>
<b>p Siterman (1997)</b>	<b>32 males</b>	<b>16-acu 16-control</b>	<b>Acupuncture</b>	<b>Sperm parameters /quality</b>	<b>P&lt;or=0.05</b>
<b>Siterman (2000)</b>	<b>40 males</b>	<b>20-acu 20-control</b>	<b>Acupuncture</b>	<b>Sperm count</b>	<b>P&lt;0.01</b>
<b>Siterman (2009)</b>	<b>57 males</b>	<b>39-acu 18-control</b>	<b>Acupuncture</b>	<b>Testes temperature/ Sperm count</b>	<b>P&lt;0.01</b>
<b>Pei (2005)</b>	<b>40 males</b>	<b>28-acu 12-control</b>	<b>Acupuncture</b>	<b>Sperm count</b>	<b>P&lt;0.05</b>
<b>Bidou (2011)</b>	<b>31 y/o male</b>	<b>NA</b>	<b>Acupuncture</b>	<b>Sperm count</b>	<b>Acupuncture is a successful method of treatment for idiopathic and non- obstructive azoospermia.</b>

### Impact of Stress on Telomere Length

Of the four articles regarding the impact of stress on telomere length, two directly linked high stress to shortened telomere length. Epel et al (2004) found that higher oxidative stress shortened telomeres in cells cultured in vivo where the continual caregiving stress of the mothers who were studied was related to telomere length. The results associate shorter telomere length with prolonged psychological stress. Zalli et al (2014) found that shortened telomeres with high telomerase activity and psychosocial factors contributed to differences in stress both psychologically and socially. The short telomere length and high telomerase activity group had fewer means of social support and higher hostility measurements. Of the two remaining articles, the results ended up being too heterogeneous to draw a conclusion.

**Table 4: Overview of Studies (Stress and Telomere Length)**

<b>1<sup>st</sup> author(date)</b>	<b>Subjects</b>	<b>Methods</b>	<b>Studied</b>	<b>Measured</b>	<b>P-value and/or Results</b>
<b>Epel (2004)</b>	<b>19-control 39- caregiving</b>	<b>Questionnaire / PBMC's</b>	<b>Telomere length</b>	<b>Stress</b>	<b><i>P</i> &lt; 0.045</b>
<b>Zalli (2014)</b>	<b>333-split into three</b>	<b>Questionnaire /stress</b>	<b>Shortened telomeres</b>	<b>Stress</b>	<b><i>P</i> = 0.046</b>

	<b>median groups</b>	<b>testing/ PBMC's Cortisol-blood &amp; saliva</b>			
--	--------------------------	--	--	--	--

### **Impact of Stress on Male Fertility**

Out of the five articles reviewed that addressed stress and male fertility, four demonstrated a correlation between these two factors, and anxiety and depression were factored under the umbrella of stress. Pilkington et al (2007) reported promising results in the management of situational and generalized anxiety using acupuncture. In 2010 they performed another study showing how depression and anxiety together constitute a significant contribution to the global burden of disease. The study found it was not possible to accurately assess the effectiveness of acupuncture for these conditions or the relative effectiveness of different treatment regimens. Studies have assessed changes in levels of neurotransmitters with some data to back this up. Errington et al (2011) found that statistically significant effects directly attributable to an acupuncture treatment lends weight to the use of acupuncture to significantly reduce the symptoms of anxiety disorders, using both human and animal subjects. Wang et al (2014) found that acupuncture improved behavior changes in rats. The objective of the study was to observe the impacts of acupuncture on various hormones and acupuncture was seen to have regulated the stress reaction. Sparrow & Golianu (2014) looked into how acupuncture may reduce stress over time. The results showed that patients tended to have an increase in their HRV(heart rate variability), a measure of physiologic stress, during treatment, after needling,

and, in some instances, an increase in HRV over weeks to months. These results indicated a relative decrease in their physiologic stress following acupuncture over the next few months.

**Table 6: Overview of Studies: (Stress[Anxiety, Depression] and Acupuncture)**

<b>1<sup>st</sup> author(date)</b>	<b>Research</b>	<b>Study/ Control</b>	<b>Measured</b>	<b>Intervention</b>	<b>Results</b>
Pilkington 2007	10 randomiz- ed control studies	4- experimental 6-control	Anxiety	Acupuncture	Insufficient evidence from research on acupuncture in the treatment of specific anxiety disorders for firm conclusions to be drawn.
Pilkington 2010	Clinical review	NA	*Anxiety *Depressi -on	Acupuncture	Because of the heterogeneity of these studies in terms of the acupuncture interventions and controls used, comparative

					effects cannot be effectively assessed
Errington 2011	Literature review	NA	Anxiety	Acupuncture	Limited/inconsistent, evidence-based recommendations regarding the adequate dose required for acupuncture to have a therapeutic effect.
Wang 2014	Male Sprague-Dawley rats	58 male rats- 5 groups randomized- varying acupuncture	Stress	Acupuncture	Acupuncture regulates stress response in rats
Sparrow (2014)	Case study	NA	Stress and heart rate variability	Acupuncture	Acupuncture reduces stress by increases heart rate variability

## **Chapter 5: Discussion**

### **Summary of findings**

The objective of this review was to review current literature related to the author's hypothesis that acupuncture increases telomere length, which results in higher sperm quality. Promising data from Omura's study in both 1998 and 2007 showed that telomere length was significantly lengthened when using acupuncture on humans. Lin in 2015 found telomere length had a correlation with male fertility, namely sperm quality. Chronic stress was studied as a strong variable that could affect telomere length and sperm quality. However, not much data was found on this specific correlation.

### **Study limitations**

Initially, the author intended to analyze available research on acupuncture's affect on telomere length, acupuncture's affect on sperm quality and telomere length's affect on sperm quality. It soon became clear that there was not enough evidence on the effect of telomere length on sperm quality, nor on the effect of acupuncture on telomeres. In addition, the author realized that the ability of acupuncture to reduce stress may play a key role in the affect acupuncture has on fertility.

Most current research found on telomere length involves the process of how it affects aging. Also, very few articles could be gathered to correlate acupuncture's affects on telomere length. When finding articles on how acupuncture can affect stress, not much research was found for stress in itself, but more on anxiety and even then not much research was available.

Other studies were found relating to this review, but they included other variables, such as acupuncture and moxabustion. They were not included because they did not meet the inclusion/exclusion criteria.

### **Theory of implication**

A large implication of theory is how the Western doctors, especially reproductive endocrinologists, can use acupuncture as an adjunct to treatment. This knowledge also empowers patients who are going through this tedious and often times emotionally heartbreaking experience. Woman many times feel depressed that they can't get pregnant and think there is something wrong with them. Since 33% of infertility is due to the male factor, the public should be made more aware of the many fertility treatment options, including less invasive approaches. This could have the potential to take a large burden off females, who statistically are the ones who receive the majority of fertility and acupuncture treatments.

### **Recommendations for additional studies**

With the current studies of epigenetics and telomere length on the forefront (Niajou, 2009) traditional Chinese medicine and namely acupuncture need to be further investigated. Research on heightened stress depleting the immune system has shown that illness comes about when the body is not in homeostasis (Saretzki & Zglinicki, 2002). As TCM practitioners, we see most patients have a large decrease in psychological stress with acupuncture. Since there have been many studies on meditation's effects on telomere length as well as meditation's effects on stress level, (Specia et al., 2000), the author suggests that further studies on acupuncture's effects on telomere length should be done. In my practice treating patients as well as my own

personal experiencing receiving acupuncture, I have observed that the body maintains homeostasis after acupuncture just as it would through a deep relaxing mediation.

### **Conclusion**

This synthesis illustrated that acupuncture can improve sperm count. These enhancements in sperm parameters could possibly give rise to improved fertility outcomes including better embryo quality, increased pregnancy rates and higher live birth rates. Evidence was also found correlating decreased telomere length with increased male infertility. This research suggests that telomere length can be increased, by decreasing levels of chronic stress. Acupuncture has been shown to decrease chronic stress, although more studies need to be done to strengthen this idea. Further controlled studies need to be done to address specific male factor infertility such as sperm count, motility, morphology and DNA fragmentation.

### **References**

- Agarwal, A., Mulgund, A., Hamada, A., Chyatte, M. (2015). A unique view on male infertility around the globe. *Reprod Biol Endocrinol.*, 13, 37.
- Aitken, R.J., Smith, T.B., Jobling, M.S., Baker, M.A., De luliis, G. N. (2014). Oxidative stress and male reproductive health. *Asian Journal of Andrology*, 16(1), 31-38.
- Allsopp R.C., Harley C.B. (1995). Evidence for a critical telomere length in senescent human fibroblasts. *Exp Cell Res.*, 219, 130–136.

- Alter, B.P., Baerlocher, G.M., Savage, S.A., Giri, N., Lansdorp, P.M. (2007). Very short telomere length by flow fluorescence in situ hybridization identifies patients with dyskeratosis congenital. *Blood*, *110*, 1439-47.
- Atzmon, G., Cho, M., Cawthon, R.M., Budagov, T., Katz, M., Yang, X. (2010). Evolution in health and medicine Sackler colloquium: genetic variation in human telomerase is associated with telomere length in Ashkenazi centenarians. *Proc Natl Acad Sci USA*, *107*(1), 1710–7.
- Aubert, G.1., Hills, M., Lansdorp, P.M. (2012). Telomere length measurement-caveats and a critical assessment of the available technologies and tools. *Mutat Res.*, *730*(1-2), 59-67.
- Baird, D.M., Britt-Compton, B., Rowson, J., Amso, N.N., Gregory, L., Kipling, D. (2006). Telomere instability in the male germline. *Hum Mol Genet.*, *15*, 45–51.
- Bass, H.W., Riera-Lizarazu, O., Ananiev, E.V., Bordoli, S.J., Rines, H.W., Phillips, R.L. (2000). Evidence for the coincident initiation of homolog pairing and synapsis during the telomere-clustering (bouquet) stage of meiotic prophase. *J Cell Sci.*, *113*, 1033–42.
- Benavides-Piccione, R and DeFelipe, J. (2003). Different Populations of Tyrosine-hydroxylase-Immunoreactive Neurons Defined by Differential Expression of Nitric Oxide Synthase in the Human Temporal Cortex *Cereb. Cortex*, *13* (3), 297-307.
- Betts, D.H., Perrault, S., Harrington, L., King, W.A. (2006). Quantitative analysis of telomerase activity and telomere length in domestic animal clones. *Methods Mol Biol.*, *325*, 149-80.
- Bidouee, F., Shamsa, A., Jalali, M. (2011). Effect of acupuncture on azoospermic male. *Saudi J Kidney Dis Transpl.*, *22*, 1039-40.
- Blackburn, E.H. (1991). Structure and function of telomeres. *Nature*, *350*, 569–573.

- Blackburn, E.H., Greider, C.W., Szostak, J.W. (2006). Telomeres and telomerase: the path from maize, Tetrahymena and yeast to human cancer and aging. *Nat Med.*, 12, 1133-38.
- Bonache, S., Mata, A., Ramos, M.D., Bassas, L., Larriba, S. (2012). Sperm gene expression profile is related to pregnancy rate after insemination and is predictive of low fecundity in normozoospermic men. *Hum Reprod.*, 27, 1556–67.
- Calado, R.T., Young, N.S. (2009). Telomere diseases. *N Engl J Med.*, 361, 2353–2365.
- Calado, R.T., Yewdell, W.T., Wilkerson, K.L., Regal, J.A., Kajigaya, S., Stratakis, C.A., Young, N.S. (2009). Sex hormones, acting on the TERT gene, increase telomerase activity in human primary hematopoietic cells. *Blood*, 114, 2236-2243.
- Carrell, D.T., Liu, L., Peterson, M. Jones, K.P., Hatasaka, H., Erickson, L., Campbell, B. (2003). Sperm DNA Fragmentation is increased in couples with unexplained recurrent pregnancy loss. *Archives of Andrology*, 49(1), 49-55.
- Chan, S.R., Blackburn, E.H. (2004). Telomeres and telomerase. *Philos Trans R Soc Lond B Biol Sci.*, 359, 109–121.
- Chao, D.M., Shen, L.L., Tjen-Alooi, S., Pitsillides, K.F., Li, P., Longhurst, J.C. (1999). Naloxone reverses inhibitory effect of electroacupuncture on sympathetic cardiovascular reflex responses. *Am J Physiol Heart Circ Physiol.*, 276, 2127–2134.
- Chen, A. (1992). An introduction to sequential electric acupuncture (SEA) in the treatment of stress related physical and mental disorders. *Acupunct Electrother Res.*, 8, 273–283.
- Chen, A., Shen, A., Li, R., Xia, Z. (2011). Effect of acupuncture-moxibustion therapy on sperm quality in infertility patients with sperm abnormality. *Journal of Acupuncture and Tuina Science*, 9(4), 219-222.

- Chen, W., Kimura, M., Kim, S., Cao, X., Srinivasan, S.R., Berenson, G.S. (2011). Longitudinal versus cross-sectional evaluations of leukocyte telomere length dynamics: age-dependent telomere shortening is the rule. *J Gerontol A Biol Sci Med Sci.*, 66, 312–9.
- Collodel, G.I., Moretti, E., Fontani, V., Rinaldi, S., Aravagli, L., Saragò, G., Capitani, S., Anichini, C. (2008). Effect of emotional stress on sperm quality. *Indian J Med Res.*, 128(3), 254-61.
- De Kretser, D.M. (1997). Male infertility. *Lancet*, 349,787–90.
- De Lange, T. (2009). How telomeres solve the end-protection problem. *Science*, 326, 948–52.
- Desai, N., Edmund, S., Taesoo, K., Ashok, A. (2010). Free Radical Theory of Aging: Implications in Male Infertility. *Urology*, 75(1), 14–19.
- Dieterle, S., Li, C., Greb, R., Bartzsch, F., Hatzmann, W., Huang, D. (2009). A prospective randomized placebo-controlled study of the effect of acupuncture in infertile patients with severe oligoasthenozoospermia. *Fertil Steril.*, 92 (4), 1340–3.
- Dohle, G.R., Weidner, W., Jungwirth, A., Colpi, G., Papp, G., Pomerol, J., Hargreave, T.B. (2011). Guidelines on Male Infertility. *European Association of Urology*, 62(2), 324-332.
- Dohle, G.R., Colpi, G., Hargreave, T.B., Papp, G., Jungwirth, A., Weidner, W. (2005). EAU Working Group on Male Infertility Guidelines. *Eur Urol.*, 48(5), 703-11.
- Edwards, R.G., Bishop, C. E. (1997). On the origin and frequency of Y chromosome deletions responsible for severe male infertility. *Molecular Human Reproduction*, 3(7) 549–554.
- Eisenberg, D.T. (2011). An evolutionary review of human telomere biology: the thrifty telomere hypothesis and notes on potential adaptive paternal effects. *Am J Hum Biol.*, 23, 149–67.
- Emery, B.R., Carrell, D.T. (2006). The effect of epigenetic sperm abnormalities on early embryo- genesis. *Asian J Androl.*, 8 (2), 131–142.

- Epel, E.S., Blackburn, E.H., Lin, J., Dhabhar, F.S., Adler, N.E., Morrow, J.D. (2004). Accelerated telomere shortening in response to life stress. *Proc Natl Acad Sci USA*, 101, 17312–17315.
- Errington-Evans, N. (2012). Acupuncture for Anxiety. *CNS Neuroscience & Therapeutics*, 18, 277–284.
- Eshkevari, L., Egan, R., Phillips, D., Tilam, J., Carney, E., Azzam, N., Amri, H., Mulroney, S.E. (2012). Acupuncture at ST36 prevents chronic stress-induced increases in neuropeptide Y in rat. *Exp Biol Med (Maywood)*, 237(1), 18-23.
- Esteves, S. A. (2013). Clinical appraisal of the genetic basis in unexplained male infertility. *J Hum Reprod Sci.*, 6(3), 176–182.
- Evenson, D., Kjersten, L., Larson, L., Jost, K. (2002). Sperm Chromatin Structure Assay: Its Clinical Use for Detecting Sperm DNA Fragmentation in Male Infertility and Comparisons With Other Techniques. *Journal of Andrology*, 23(1), 25–43.
- Eshkevari L., Egan, R., Phillips, D., Tilan, J., Carney, E., Azzam, N., Amri, H., Mulroney, S.E. (2012). Acupuncture at ST36 prevents chronic stress-induced increases in neuropeptide Y in rat. *Exp Biol Med (Maywood)*, 237(1), 18-23.
- Eshkevari, L., Mulroney, S.E., Egan, R., Lao, L. (2015). Effects of Acupuncture, RU-486 on the Hypothalamic-Pituitary-Adrenal Axis in Chronically Stressed Adult Male Rats. *Endocrinology*, 156(10), 3649-60.
- Eshkevari, L., Permaul, E., Mulroney, S.E. (2013). Acupuncture blocks cold stress-induced increases in the hypothalamus-pituitary-adrenal axis in the rat. *J Endocrinol*, 217(1), 95-104.

- Fajkus, J., Dvoráková, M., Sýkorová, E. (2008). Analysis of telomeres and telomerase. *Methods Mol Biol.*, 463, 267-96.
- Ferlin, A., Raicu, F., Gatta, V., Zuccarello, D., Palka, G. (2007). Male infertility: role of genetic background. *Reprod Biomed.*, 14, 734–45.
- Ferlin, A. (2012). New genetic markers for male fertility. *Asian J Androl.*, 14(6), 807–808.
- Flores, I., Blasco, M.A. (2010). The role of telomeres and telomerase in stem cell aging. *FEBS Lett.*, 584, 3826–30.
- Foresta, C. (2007). Male infertility: role of genetic background. *Reprod Biomed Online*, 14, 734–45.
- Gatta, V., Raicu, F., Ferlin, A., Antonucci, I., Scioletti, A.P. (2010). Testis transcriptome analysis in male infertility: new insight on the pathogenesis of oligo-azoospermia in cases with and without AZFc microdeletion. *BMC Genomics*, 11, 401.
- Gnoth, C., Godehardt, E., Frank-Herrmann, P., Friol, K., Tigges, J., Freundl, G. (2005). Definition and prevalence of subfertility and infertility. *Hum Reprod.*, 20(5), 1144-7.
- Gui, Z., Zhang, Z., Wang, G., Han, F., Han, L., Wang, K., Liu, J., Li, W. (2011). Effects of Electroacupuncture Pretreatment on Inflammatory Response and Acute Kidney Injury in Endotoxaemic Rats. *The Journal of International Medical research*, 39, 1783-1797.
- Gurfinkel, E., Cedenho, A.P., Yamamura, Y., Srougi, M. (2003). Effects of acupuncture and moxa treatment in patients with semen abnormalities. *Asian J Androl.*, 5(4), 345-8.
- Hanna, C.W., Bretherick, K.L., Gair, J.L., Fluker, M.R., Stephenson, M.D., Robinson, W.P. (2009). Telomere length and reproductive aging. *Hum Reprod.*, 24, 1206–11.
- Harley, C.B., Futcher, A.B., Greider, C.W. (1990). Telomeres shorten during aging of human fibroblasts. *Nature*, 345, 458–460.

- Heiss, N.S. (1998). X-linked dyskeratosis congenita is caused by mutations in a highly conserved gene with putative nucleolar functions. *Nat Genet*, *19*, 32-38.
- Hemann, M.T., Strong, M.A., Hao, L.Y., Greider, C.W. (2001). The shortest telomere, not average telomere length, is critical for cell viability and chromosome stability. *Cell*, *107*(1), 67-77.
- Herrera, E., Samper, E., □Martín-Caballero, J., Flores, J.M., Lee, H.W., Blasco, M.A. (1999). Disease states associated with telomerase deficiency appear earlier in mice with short telomeres. *The EMBO Journal*, *18*(11), 2950–2960.
- Houben, J.M. (2011). Telomere length and mortality in elderly men: the Zutphen Elderly Study. *J Gerontol A Biol Sci Med Sci.*, *66*, 38- 44.
- Huddleston, H.G., Cedars, M.I., Sohn, S.H., Giudice, L.C., Fujimoto, V.Y. (2010). Racial and ethnic disparities in reproductive endocrinology and infertility. *Am J Obstet Gynecol.*, *202*(5), 413-9.
- Inoue, K., Aikawa, H., Yamauchi, T. (2003). Autonomic nervous activity and arousal level of patients with anxiety disorder. *Clin Neurophysiol.*, *8*, 499–502.
- Jain ,T. (2006). Socioeconomic and racial disparities among infertility patients seeking care. *Fertil Steril.*, *85*(4), 876-81.
- Jiao Mu, J., Wei, L.X. (2002). Telomere and telomerase in oncology. *Cell Research*, *12*, 1–7.
- Kalmbach, K., Fontes-Antunes, D.M., Dracxler, R.C., Knier, T.W., Seth-Smith, M.L., Wang, F., Liu, L., Keefe, D.L. (2013). Telomeres and human reproduction. *Fertil Steril.*, *99*(1), 23-29.
- Kim, N.W., Piatyszek, M.A., Prowse, K.R. (1994). Specific association of human telomerase activity with immortal cells and cancer. *Science*, *266*, 2011–5.

- Keefe, D.L., Liu L. (2009). Telomeres and reproductive aging. *Reprod Fertil Dev.*, 21, 10–4.
- Kimura, M., Cherkas, L.F., Kato, B.S., Demissie, S., Hjelmborg, J.B., Brimacombe, M. (2008). Offspring's leukocyte telomere length, paternal age, and telomere elongation in sperm. *PLoS Genet.* 4(37).
- Knardahl, S., Elam, M., Olausson, B., Wallin, B.G. (1998). Sympathetic nerve activity after acupuncture in humans. *Pain*, 8, 19–25.
- Kondo, T., Kawamoto, M. (2014). Acupuncture and moxibustion for stress-related disorders. *Biopsychosoc Med.*, 8, 7.
- Kosova, G, Scott, N.M., Niederberger, C., Prins, G.S., Ober, C. (2012). Genome-wide association study identifies candidate genes for male fertility traits in humans. *Am J Hum Genet.* 90, 950–61.
- Lee, J.S., Jeong, S.W., Cho, S.W., Juhn, J.P., Kim K.W. (2001). Relationship between Initial Telomere Length, Initial Telomerase Activity, Age, and Replicative Capacity of Nucleus Pulposus Chondrocytes in Human Intervertebral Discs: What Is a Predictor of Replicative Potential? Switching and signaling at the telomere. *Cell*, 106, 661–673.
- Lin, D., Wu, Q., Lin., Borlongan, C.V., He, Z., Tan, J., Cao, C., Zhou, S. (2015). Brain-derived Neurotrophic Factor Signaling Pathway: Modulation by Acupuncture in Telomerase Knockout Mice. *Alternative Therapies* 21(6), 36.
- Liu, L., Bailey, S.M., Okuka, M., Munoz, P., Li, C., Zhou, L. (2007). Telomere lengthening early in development. *Nat Cell Biol.*, 9, 1436–41.
- Liu, J., Huang, H., Xu, X., Chen, J.D. (2012). Effects and possible mechanisms of acupuncture at ST36 on upper and lower abdominal symptoms induced by rectal distension in healthy volunteers. *Am J Physiol Regul Integr Comp Physiol.*, 8, 209–217.

- Liu, L., Trimarchi, J.R., Smith, P.J., Keefe, D.L. (2002). Mitochondrial dysfunction leads to telomere attrition and genomic instability. *Aging Cell*, 1, 40–6.
- Lopez, G., Lafuente, R., Checa, M., Carreras, R., Brassesco, M. (2013). Diagnostic value of sperm DNA fragmentation and sperm high-magnification for predicting outcome of assisted reproduction treatment. *Asian J Androl.*, 15(6), 790-794.
- Machev, N.1., Gosset, P., Viville, S. (2005). Chromosome abnormalities in sperm from infertile men with normal somatic karyotypes: teratozoospermia. *Cytogenet Genome Res.*, 111(3-4), 352-7.
- Mayorga-Torres, B.J., Camargo, M., Cadavid, Á.P., du Plessis, S.S., Cardona Maya, W.D. (2016). Are oxidative stress markers associated with unexplained male infertility?. *Andrologia*.
- McGrady, A. V. (1984). Effects of Psychological Stress on Male Reproduction: A Review. *Archives of Andrology*, 13(1).
- Middlekauff, H., Yu, J.L., Hui, K. (2001). Acupuncture effects on reflex responses to mental stress in humans. *American Journal of Physiology - Regulatory, Integrative and Comparative Physiology*, 280(5), 1462-1468.
- Miller, D., Ostermeier, G.C. (2006). Towards a better understanding of RNA carriage by ejaculate spermatozoa. *Hum Reprod Update*, 12, 757–67.
- Minagawa, M., Kurono, Y., Ishigami, T., Yamada, A., Kakamu, T., Akai, R. (2013). Site-specific organ-selective effect of epifascial acupuncture on cardiac and gastric autonomic functions. *Auton Neurosci.*, 8, 151–154.
- Montpetit, A.J., Alhareeri, A.A., Montpetit, M., Starkweather, A.R., Elmore, L.W., Filler, K.,

- Mohanraj, L., Burton, C.W., Menzies, V.S., Lyon, D.E., Jackson-Cook, C.K. (2014).  
Telomere length: a review of methods for measurement. *Nurs Res.*, 63(4), 289-99.
- Moskovtsev, S.I., Willis, J., White, J., Mullen, J.B. (2010). Disruption of telomere-telomere interactions associated with DNA damage in human spermatozoa. *Systems biology in reproductive medicine*, 56, 407–12.
- Mu, J., Wei, L.X. (2002). Telomere and telomerase in oncology. *Cell Research*. 12, 1–7.
- Nakamura, M., Yukawa, S. (2010). The importance of the counseling aspects of acupuncture/moxibustion treatments. *Jpn J Couns Sci.*, 8, 192–201.
- Nargund, V.H .Effects of psychological stress on male fertility. (2015). *Nat Rev Urol.*, 12(7), 373-82.
- Neto, F.T., Bach, P.V., Najari, B.B., Li, P.S., Goldstein, M. (2016). Genetics of Male Infertility. *Curr Urol Rep.*, 17(10), 70.
- Ngai, S.P., Jones, A.Y. (2013). Changes in skin impedance and heart rate variability with application of Acu-TENS to BL 13 (feishu). *J Altern Complement Med.*, 8, 558–563.
- Niederberger, C . (2016). DNA Fragmentation in Brighter Sperm Predicts Male Fertility Independently from Age and Semen Parameters. *J Urol.*, 196(2), 520-1.
- Njajou, O.T. (2009). Association between telomere length, specific causes of death, and years of healthy life in health, aging, and body composition, a population-based cohort study. *J Gerontol A Biol Sci Med Sci.*, 64, 860-64.
- Ober, C., Hyslop, T., Hauck, W.W. (1999). Inbreeding effects on fertility in humans: evidence for reproductive compensation. *Am J Hum Genet*, 64, 225–31.

O'Flynn O'Brien, K.L., Varghese, A.C., Agarwal, A. (2010). The genetic causes of male factor infertility: A review. *Fert Steril.*, 93(1), 1–12.

Omura, Y., Chen, Y., Lermann, O., Jones, M., Duvvi, H., Shimotsura, Y. (2010). Effects of Transcutaneous Electrical Stimulation (1 pulse/sec) Through Custom-made Disposable Surface Electrodes Covering Omura's ST36 Area of Both Legs on Normal Cell Telomeres, Oncogen C-fosAb2, Integrin  $\alpha 5\beta 1$ , Chlamydia Trachomatis, etc. in Breast Cancer & Alzheimer Patients. *Acupuncture & Electro-Therapeutics Research*, 35(3-4), 147-185.

Omura, Y., Chen, Y., Lu, D.P., Shimotsura, Y., Ohki, M., Duvvi, H. (2007). Anatomical Relationship Between Traditional Acupuncture Point ST 36 and Omura's ST 36 (true ST 36) With Their Therapeutic Effects: 1) Inhibition of Cancer Cell Division by Markedly Lowering Cancer Cell Telomere While Increasing Normal Cell Telomere, 2) Improving Circulatory Disturbances, with Reduction of Abnormal Increase in High Triglyceride, L-Homocystein, CRP, or Cardiac Troponin I & T in Blood By the Stimulation of Omura's ST 36 - - - Part 1. *Acupuncture & Electro-Therapeutics Research*, 32(1-2), 31-70.

Omura, Y., Shimotsura, Y., Ooki, M., Noguchi, T. (1998). Estimation of the Amount of Telomere Molecules in Different Human Age Groups and the Telomere Increasing Effect of Acupuncture and Shiatsu on St.36, Using Synthesized Basic Units of the Human Telomere Molecules as Reference Control Substances for the Bi-Digital O-Ring Test Resonance Phenomenon. *Acupuncture & Electro-Therapeutics Research*, 23(3-4), 185-206.

- Owen, C.M., Goldstein, E.H., Clayton, J.A. Segars, J.H. (2013). Racial and ethnic health disparities in reproductive medicine: an evidence-based overview. *Semin Reprod Med.*, 31(5), 317-24.
- Park, H.J., Kim, H.Y., Hahm, D.H., Lee, H., Kim, K.S., Shim, I. (2010). Electroacupuncture to ST36 ameliorates behavioral and biochemical responses to restraint stress in rats. *Neurol Res.*, 8(1), 111–115.
- Pei, J., Strehler, E., Noss, U., Abt, M., Piomboni, P., Baccetti, B., Sterzik, K. (2005). Quantitative evaluation of spermatozoa ultrastructure after acupuncture treatment for idiopathic male infertility. *Fertil Steril.*, 84(1), 141-7.
- Pickering, C.M., Byrne, J. (2013). The benefits of publishing systematic quantitative literature reviews for PhD candidates and other early career researchers. *Higher Education Research and Development*, 33(3), 534-548.
- Pilkington, K. (2010). Anxiety, depression and acupuncture: A review of the clinical research. *Auton Neurosci.*, 157(1-2), 91-5
- Pilkington, K., Kirkwood, G., Rampes, H., Cummings, M., Richardson, J. (2007). Acupuncture for anxiety and anxiety disorders – a systematic literature review. *Acupunct Med.*, 25, 1-10.
- Pomeranz, B., Bibic, L. (1988) Naltrexone, an opiate antagonist, prevents but does not reverse the analgesia produced by electroacupuncture. *Brain Res.*, 452, 227–231.
- Reynolds, R.M. (2016). Stress and the Brain: from Fertility to Senility. *J Neuroendocrinol.*, 28, 8.
- Rode, L., Nordestgaard, B.G., Bojesen, S.E. (2016). Long telomeres and cancer risk among 95 568 individuals from the general population. *Int J Epidemiol.*

- Rodríguez, S., Goyanes, V., Segrelles, E., Blasco, M., Gosálvez, J., Fernández, J.L. (2005). Critically short telomeres are associated with sperm DNA fragmentation. *Fertil Steril.*, *84(4)*, 843-5.
- Rousseaux, S., Caron, C., Govin, J., Lestrat, C., Faure, A., Khochbin, S. (2005). Establishment of male-specific epigenetic information. *Gene*, *345*, 139–153.
- Rudolph, K.L. (2007). Telomere Shortening and Telomerase Activation during Cancer Formation. In K. Lenhard Rudolph (Ed.), *Telomeres and Telomerase in Aging, Disease and Cancer: Molecular Mechanisms of Adult Stem Cell Aging* (pp.223-227). Ulm, Germany: Springer Science & Business Media.
- Saretzki, G., Von Zglinicki, T. (2002). Replicative aging, telomeres, and oxidative stress. *Ann N Y Acad Sci.*, *959*, 24–29.
- Schaeffer, A.J., Hotaling, J.M. (2016). Claims-based analysis of male infertility: a cautious step in the right direction. *Fertil Steril.*, *105(3)*, 599-600.
- Scheinberg, P. (2010). Association of telomere length of peripheral blood leukocytes with hematopoietic relapse, malignant transformation, and survival in severe aplastic anemia. *JAMA*, *304*, 1358-64.
- Scherthan, H. (2007). Telomere attachment and clustering during meiosis. *Cell Mol Life Sci.*, *64*, 117–24.
- Shah, K., Sivapalan, G., Gibbons, N., Tempest, H., Griffin, D.K. (2003). The genetic basis of infertility. *Reproduction*, *126(1)*, 13-25.
- Sharara, F.I. (2013). Introduction: The Scope of the Topic In F.I. Sharara (Ed.) *Ethnic Differences in Fertility and Assisted Reproduction* (pp. 1-5). Reston, VA: Springer Science & Business Media.

- Shay, J.W., Wright, W.E. (2011). Role of telomeres and telomerase in cancer. *Semin Cancer Boil.*, 21(6), 349-353.
- Shay, J.W., Wright, W.E. (2010). Telomeres and telomerase in normal and cancer stem cells. *FEBS Lett.*, 584(17), 3819–3825.
- Shay, J.W., Zou, Y., Hiyama, E., Wright, W.E. (2001). Telomerase and cancer. *Hum. Mol. Genet.*, 10(7), 677-685.
- Shea, A.K., Kamath, M.V., Fleming, A, Streiner, D.L., Redmond, K., Steiner, M. (2008). The effect of depression on heart rate variability during pregnancy. A naturalistic study. *Clin Auton Res.*, 8, 203–212.
- Sheps, M.C. (1965). An analysis of reproductive patterns in an American isolate. *Popul Stud.*, 19, 65–80.
- Sherman, S., Eltes, F., Wolfson, V., Zabludovsky, N., Bartoov, B. (1997). Effect of acupuncture on sperm parameters of males suffering from subfertility related to low sperm quality. *Arch Androl.*, 39(2), 155-61.
- Sinclair, S. (2000). Male infertility: nutritional and environmental considerations. *Altern Med Rev.*, 5(1), 28-38.
- Siterman, S., Eltes, F., Schechter, L., Maimon, Y., Lederman, H., Bartoov, B. (2009). Success of acupuncture treatment in patients with initially low sperm output is associated with a decrease in scrotal skin temperature. *Asian J Androl.*, 11(2), 200-8.
- Singh, K., Jaiswal, D. (2011). Human male infertility: a complex multifactorial phenotype. *Reprod Sci.*, 18(5), 418-25.

- Siterman, S., Eltes, F., Wolfson, V., Lederman, H., Bartoov, B. (2000). Does acupuncture treatment affect sperm density in males with very low sperm count? A pilot study. *Andrologia*, 32(1), 31-9.
- Siterman, S., Eltes, F., Wolfson, V., Zabludovsky, N., Bartoov, B. (1997). Effect of acupuncture on sperm parameters of males suffering from subfertility related to low sperm quality. *Arch Androl.*, 39(2), 155-61.
- Sparrow, K., Golianu, B. (2014). Does Acupuncture Reduce Stress Over Time? A Clinical Heart Rate Variability Study in Hypertensive Patients. *Med Acupunct.*, 26(5), 286–294.
- Specia, M., Carlson, L., Goodey, E, Angen, M. (2000). A Randomized, Wait-List Controlled Clinical Trial: The Effect of a Mindfulness Meditation-Based Stress Reduction Program on Mood and Symptoms of Stress in Cancer Outpatients. *Psychosomatic Medicine*. 62, 613–622.
- Sun, J.P., Pei, H.T., Jin, X.L., Yin, L., Tian, Q.H., Tian, S.J. (2005). Effects of acupuncturing Tsusanli (ST36) on expression of nitric oxide synthase in hypothalamus and adrenal gland in rats with cold stress ulcer. *World J Gastroenterol.*, 11, 4962–618.
- Sylvest, R., Fürbringer, J.K., Schmidt, L., Pinborg, A. (2016). Infertile men's needs and assessment of fertility care. *Ups J Med Sci.*,8, 1-7.
- Taylor, A. (2003). ABC of subfertility Extent of the problem. *Oxford Journals Medicine & Health Human Reproduction*, 20(5), 1144-1147.
- Tempest, H.G., Homa, S.T., Zhai, X.P., Griffin, D.K. (2005). Significant reduction of sperm disomy in six men: effect of traditional Chinese medicine? *Asian J Androl.*, 7(4), 419-25.
- Thilagavathi ,J., Venkatesh, S., Dada, R. (2012). Telomere length in reproduction. *Andrologia*, 45(5), 289-304.

- Tonhajzerova, I., Ondrejka, I., Javorka, K., Turianikova, Z., Farsky, I., Javorka, M. (2010). Cardiac autonomic regulation is impaired in girls with major depression. *Prog Neuropsychopharmacol Biol Psychiatry*, 8, 613–618.
- Treff, N.R., Su, J., Taylor, D., Scott, R.T. (2011). Telomere DNA deficiency is associated with development of human embryonic aneuploidy. *PLoS Genet.*, 7, 6.
- Wang, S, Zhang, J., Qie, L. (2014). Acupuncture Relieves the Excessive Excitation of Hypothalamic-Pituitary-Adrenal Cortex Axis Function and Correlates with the Regulatory Mechanism of GR, CRH, and ACTHR. *Evidence-based Compl Alt Med.*, 6.
- Weischer, M., Nordestgaard, B.G., Cawthon, R.M., Freiberg, J.J., Tybjærg-Hansen, A., Bojesen, S.E. (2013). Short telomere length, cancer survival, and cancer risk in 47102 individuals. *J Natl Cancer Inst.*, 105(7), 459-68.
- Wesch, N.L., Burlock, L.J., Gooding, R.J. (2016). Critical telomerase activity for uncontrolled cell growth. *Phys Biol.*,13(4).
- Xiao, M., Yang, X., ,Zhong, Run, Xie, Z., Li, J., Zhou, Y., Tian, N., Yuan, H., Zhao, H. (2013). Experimental study of the influence on moxibustion for telomere length in brain cell of subacute aging mice. *Beijing Journal of Traditional Chinese Medicine*, 1.
- Xu, Z., Duc, K.D., Holcman, D., Teixeira, M.T. (2013). The length of the shortest telomere as the major determinant of the onset of replicative senescence. *Genetics*, 194, 847–857.
- Yao, T., Andersson, S., Thoren, P. (1982) Long-lasting cardiovascular depression induced by acupuncture-like stimulation of the sciatic nerve in unanaesthetized spontaneously hypertensive rats. *Brain Res.*, 240, 77–85.
- Yook, T., Yu, J., Lee, H., Song, B., Kim, L., Roh, J. (2009). Comparing the effects of distilled *Rehmannia glutinosa*, Wild Ginseng and *Astragali Radix* pharmacopuncture with heart r

ate variability (HRV): a randomized, sham-controlled and double-blind clinical trial. *J Acupunct Meridian Stud.*, 8, 239–247.

Zalli, A., Carvalho, L.A., Lin, J., Hamer, M., Erusalimsky, J.D., Blackburn, E.H., Steptoea, A. (2014). Shorter telomeres with high telomerase activity are associated with raised allostatic load and impoverished psychosocial resources. *Proc Natl Acad Sci USA*, 111(12), 4519–4524.

Zhu, X., Han, W., Xue, W., Zou, Y., Xie, C., Du, J., Jina, G. (2016). The association between telomere length and cancer risk in population studies. *Sci Rep.*, 6, 22243.

**Appendix 1: Table 4**

Prognostic accuracy of sperm DNA fragmentation to predict outcome after IVF/ICSI  
(Lopez et al, 2013)

<i>Statistical characteristics</i>	<i>Pregnant</i>	<i>Non-pregnant</i>	<i>P value</i>
DNA fragmentation (%)	18.90±7.71	21.61±11.59	0.102
Area under ROC curve (95% CI)	0.546 (0.450–0.642)		
Cutoff (%)	25.5		0.029
Sensitivity (%)	86.2		
Specificity (%)	28.9		
Positive predictive value (%)	48.7		
Negative predictive value (%)	72.7		

Abbreviations: ICSI, intracytoplasmic sperm injection; IVF, *in vitro* fertilisation; ROC, receiver operating characteristic.

## Appendix 2: Table 5

### Methods Used to Assess Telomere Length (Montpetit et al., 2009)

Method	Analyte	Measures			
		Average	Chromosome-Specific	Resolution (kb)	Optimally Suited for Large Studies
TRF	DNA	Yes	No	1.0 <sup>a</sup>	No
qPCR	DNA	Yes	No	0.7 <sup>b</sup> , 0.9 <sup>c</sup> , 1.0 <sup>d</sup>	Yes
MMqPCR					
aTLqPCR					
STELA	DNA	No	Yes	0.1 <sup>a</sup>	No
Q-FISH	Metaphase chromosomes	Yes	Yes	0.15–0.3 <sup>a, b</sup>	No
	Interphase nuclei (telomere)	Yes	No	0.15–0.3 <sup>a, b</sup>	No
PRINS	Metaphase chromosomes	Yes	Yes	0.3 <sup>a</sup>	No
	Interphase nuclei (telomere)	Yes	No	0.3 <sup>a</sup>	No
Flow-FISH	Interphase nuclei	Yes	No	0.2–0.3 <sup>a</sup>	No
HT Q-FISH	Interphase nuclei	Yes	No	0.2–0.3 <sup>b</sup>	Yes

*Note.* aTL = Absolute telomere length. DNA = Deoxyribonucleic acid. HT Q-FISH = High throughput quantitative fluorescence in situ hybridization. kb = kilobase. MMqPCR = Monochrome multiplex quantitative polymerase chain reaction. qPCR = Quantitative polymerase chain reaction. STELA = Single Telomere Length Analysis, Universal STELA. TRF = Terminal restriction fragment. PRINS = Primed in situ subtype of Q-FISH. Q-FISH = Quantitative fluorescence in situ hybridization.

**Appendix 3: IRB Proposal**

## Appendix 4: Curriculum Vitae

### EDUCATION

---

May 2015-present (anticipated graduation April 2017)  
Yo San University  
\* Doctoral candidate of Acupuncture and Oriental Medicine  
\* Reproductive Medicine specialty

September 2000-2004  
Emperor's College  
\* Master of Traditional Oriental Medicine, *Magna Cum Laude*

September 1995-December 1998  
University of Florida  
\* Bachelor of Science, Psychology, *Honors*  
\* Pre-medicine full course work  
\* Business, Minor

### PROFESSIONAL EXPERIENCE

---

December 2004-present  
Romy Simone: Cosmetic & Medical Acupuncture, Santa Monica, CA  
Acupuncturist

September 2012-December 2013  
Chinese Healing Arts Center, Santa Monica, CA  
Independent Contractor-Cosmetic Acupuncture Specialty

March 2010-September 2012  
Gentle Journey Spa, Santa Monica, CA  
Independent Contractor-Cosmetic Acupuncture Specialty

February 2008-March 2010  
Tola Wellness Spa, Malibu, CA  
Independent Contractor-Cosmetic Acupuncture Specialty

December 2006-February 2008

Piel Skincare Spa, Beverly Hills, CA  
Independent Contractor-Cosmetic Acupuncture Specialty

December 2004-December 2006  
Veronica Skincare Spa, Malibu, CA  
Independent Contractor-Cosmetic Acupuncture Specialty

## LICENSES

---

- \* State of California Licensed Acupuncturist
- \* NCCAOM Accredited
- \* ACE Fitness Licensed

## TEACHING EXPERIENCE

---

May 2015-present  
Yo San University, Los Angeles, CA Qi Gong Instructor

October 2005-present  
Cosmetic Acupuncture Seminars, Santa Monica, CA  
Private one-on-one cosmetic acupuncture training

July 2006-present  
Various conferences and presentations, Los Angeles, CA Guest Lecturer

January 2005-2013  
Private lecture series on health and anti-aging, Malibu, CA Sole Lecturer

September-December 2013  
Emperor's College, Santa Monica, CA Assistant Patient Supervisor to Dr. Kim

September-December 2013  
Emperor's College, Santa Monica, CA Assistant Herb Dispensary Supervisor

August 2012  
Emperor's College, Santa Monica, CA Assistant Clinic Supervisor

September-December 2008  
Emperor's College, Santa Monica, CA Medical Qi Gong Instructor

September 2004  
UCLA, Los Angeles, CA Medical Qi Gong Instructor

September 2002-2004  
Emperor's College, Santa Monica, CA Medical Qi Gong Assistant Instructor

#### SPECIAL INTERESTS

---

Charity Sponsor:

January 2012-present  
Unlikely Heroes, Los Angeles, CA

February 2013-present  
Face Forward, Beverly Hills, CA

January 2014-present  
Kevin Richardson, PAW Conservation Trust, Northeast South Africa

January 2014-present  
The Jane Goodall Institute, South Africa

#### HOBBIES, INTERESTS AND ACTIVITIES

---

Argentine Tango, Gyrotonics, Show jumping-Horses, Electric Guitar, Surfing, Painting, Sculpture, Meditation, Yoga, Swimming, Ecstatic Dance, Radical Honesty